

On colour-preserving automorphisms of Cayley graphs

Ademir Hujdurović, Klavdija Kutnar

University of Primorska, FAMNIT, Glagoljaška 8, 6000 Koper, Slovenia

Dave Witte Morris, Joy Morris

*Department of Mathematics and Computer Science, University of Lethbridge,
Lethbridge, Alberta, T1K 3M4, Canada*

March 27, 2015

Abstract

We study the automorphisms of a Cayley graph that preserve its natural edge-colouring. More precisely, we are interested in groups G , such that every such automorphism of every connected Cayley graph on G has a very simple form: the composition of a left-translation and a group automorphism. We find classes of groups that have this property, and we determine the orders of all groups that do not have this property. We also have analogous results for automorphisms that permute the colours, rather than preserving them.

Keywords: Cayley graph, automorphism, colour-preserving, colour-permuting

Math. Subj. Class.: 05C25

Contents

1	Introduction	2
2	Examples of non-CCA groups	4
3	Direct products and semidirect products	8
4	Abelian groups	10
5	Generalized dihedral groups	12
6	Groups of odd order	14
7	Groups of small order	20
A	Dicyclic (and other) groups omitted from the proof of Proposition 7.3	25
B	Strongly CCA groups omitted from the proof of Proposition 7.3	27

E-mail addresses: ademir.hujdurovic@upr.si (Ademir Hujdurović), klavdija.kutnar@upr.si (Klavdija Kutnar), dave.morris@uleth.ca (Dave Witte Morris), joy.morris@uleth.ca (Joy Morris)

1 Introduction

Definitions 1.1. Let S be a subset of a group G , such that $S = S^{-1}$. (All groups and all graphs in this paper are finite.)

- The *Cayley graph* of G , with respect to S , is the graph $\text{Cay}(G; S)$ whose vertices are the elements of G , and with an edge $x - xs$, for each $x \in G$ and $s \in S$.
- $\text{Cay}(G; S)$ has a natural edge-colouring. Namely, each edge of the form $x - xs$ is coloured with the set $\{s, s^{-1}\}$. (In order to make the colouring well-defined, it is necessary to include s^{-1} , because $x - xs$ is the same as the edge $xs - x$, which is of the form $y - ys^{-1}$, with $y = xs$.)

Note that $\text{Cay}(G; S)$ is connected if and only if S generates G . Also note that a permutation φ of G is a colour-preserving automorphism of $\text{Cay}(G; S)$ if and only if we have $\varphi(xs) \in \{\varphi(x)s^{\pm 1}\}$, for each $x \in G$ and $s \in S$.

For each $g \in G$, the left translation $x \mapsto gx$ is a colour-preserving automorphism of $\text{Cay}(G; S)$. In addition, if α is an automorphism of the group G , such that $\alpha(s) \in \{s^{\pm 1}\}$ for all $s \in S$, then α is also a colour-preserving automorphism of $\text{Cay}(G; S)$. We will see that, in many cases, every colour-preserving automorphism of $\text{Cay}(G; S)$ is obtained by composing examples of these two obvious types.

Definitions 1.2. Let G be a group.

1. A function $\varphi: G \rightarrow G$ is said to be *affine* if it is the composition of an automorphism of G with left translation by an element of G . This means $\varphi(x) = \alpha(gx)$, for some $\alpha \in \text{Aut } G$ and $g \in G$.
2. A Cayley graph $\text{Cay}(G; S)$ is *CCA* if all of its colour-preserving automorphisms are affine functions on G . (CCA is an abbreviation for the *Cayley Colour Automorphism* property.)
3. We say that G is *CCA* if every connected Cayley graph on G is CCA.

Here are some of our main results:

Theorem 1.3.

1. There is a non-CCA group of order n if and only if $n \geq 8$ and n is divisible by either 4, 21, or a number of the form $p^q \cdot q$, where p and q are prime (see Corollary 6.13 and Remark 6.14).
2. An abelian group is not CCA if and only if it has a direct factor that is isomorphic to either $\mathbb{Z}_4 \times \mathbb{Z}_2$ or a group of the form $\mathbb{Z}_{2^k} \times \mathbb{Z}_2 \times \mathbb{Z}_2$, with $k \geq 2$ (see Proposition 4.1).
3. Every dihedral group is CCA (see Corollary 5.4).
4. No generalized dicyclic group or semidihedral group is CCA, except $\mathbb{Z}_4$ (see Corollary 2.8).
5. Every non-CCA group of odd order has a section that is isomorphic to either the nonabelian group of order 21 or a certain generalization of a wreath product (called a semi-wreathed product) (see Theorem 6.8).
6. If $G \times H$ is CCA, then G and H are both CCA (see Proposition 3.1). The converse is not always true (for example, $\mathbb{Z}_4 \times \mathbb{Z}_2$ is not CCA), but it does hold if $\gcd(|G|, |H|) = 1$ (see Proposition 3.2).

We also consider automorphisms of $\text{Cay}(G; S)$ that permute the colours, rather than preserving them:

Definitions 1.4.

- An automorphism α of a Cayley graph $\text{Cay}(G; S)$ is *colour-permuting* if it respects the colour classes; that is, if two edges have the same colour, then their images under α must also have the same colour. This means there is a permutation π of S , such that $\alpha(gs) \in \{\alpha(g)\pi(s)^{\pm 1}\}$ for all $g \in G$ and $s \in S$ (and $\pi(s^{-1}) = \pi(s)^{-1}$).
- A Cayley graph $\text{Cay}(G; S)$ is *strongly CCA* if all of its colour-permuting automorphisms are affine functions on G .
- We say that G is *strongly CCA* if every connected Cayley graph on G is strongly CCA.

Note that every strongly CCA group is CCA, since colour-preserving automorphisms are colour-permuting (with π being the identity map on S). The converse is not true. For example, every dihedral group is CCA (as was mentioned above), but it is not strongly CCA if its order is of the form $8k + 4$ (see Proposition 5.6). However, the converse does hold for at least two natural families of groups:

Theorem 1.5. A CCA group is strongly CCA if either:

1. it is abelian (see Proposition 4.1), or
2. it has odd order (see Proposition 6.4).

Remarks 1.6.

1. It follows from Theorems 1.3(2) and 1.5(1) that every cyclic group is strongly CCA. This is also a consequence of the main theorem of [10].
2. Groups of even order seem far more likely to fail to be strongly CCA than groups of odd order. For example, of the 28 groups of order less than 32 that are not strongly CCA, only one has odd order (see Section 7). In fact, there are only three groups of odd order less than 100 that are not strongly CCA: the non-abelian group G_{21} of order 21, the group $G_{21} \times \mathbb{Z}_3$ of order 63, and the wreath product $\mathbb{Z}_3 \wr \mathbb{Z}_3$, which has order 81 (see Proposition 7.4).
3. If the subgroup consisting of all left-translations is normal in the automorphism group of the Cayley graph $\text{Cay}(G; S)$, then $\text{Cay}(G; S)$ is said to be *normal* [13]. It is not difficult to see that every normal Cayley graph is strongly CCA (cf. Remark 6.2), and that every automorphism of a normal Cayley graph is colour-permuting.
4. The notion of (strongly) CCA generalizes in a natural way to the setting of Cayley digraphs $\overrightarrow{\text{Cay}}(G; S)$, by putting the colour s on each directed edge of the form $x \rightarrow xs$. (There is no need to include s^{-1} in the colour.) However, it is very easy to see that if $\overrightarrow{\text{Cay}}(G; S)$ is connected, then every colour-preserving automorphism of $\overrightarrow{\text{Cay}}(G; S)$ is left-translation by some element of G [12, Thm. 4-8, p. 25], and that every colour-permuting automorphism is affine [3, Lem. 2.1]. Therefore, both notions are completely trivial in the directed setting. However, there has been some interest in determining when every automorphism of $\overrightarrow{\text{Cay}}(G; S)$ is colour-permuting [1, 2] (in which case, the Cayley digraph is normal, in the sense of (3)).

Acknowledgments. We thank an anonymous referee for numerous helpful comments that improved the exposition.

D. W. M. and J. M. thank the Faculty of Mathematics, Natural Sciences and Information Technologies of the University of Primorska (Slovenia) for its hospitality during the visit that gave rise to this research project.

The work of A. H. was partially supported by research program P1-0285 from the Slovenian Research Agency. The work of K. K. was partially supported by research program P1-0285 and research projects N1-0011, J1-6743, and J1-6720 from the Slovenian Research Agency. The work of J. M. was partially supported by a research grant from the Natural Sciences and Engineering Research Council of Canada.

2 Examples of non-CCA groups

Remark 2.1. Since automorphisms are the only affine functions that fix the identity element e (and left-translations are colour-preserving automorphisms of any Cayley graph), it is easy to see that if $\text{Cay}(G; S)$ is CCA, then every colour-preserving automorphism that fixes the identity is an automorphism of the group G . More precisely:

A Cayley graph $\text{Cay}(G; S)$ is CCA if and only if, for every colour-preserving automorphism φ of $\text{Cay}(G; S)$, such that $\varphi(e) = e$, we have $\varphi \in \text{Aut } G$.

The same is true with “strongly CCA” in the place of “CCA,” if “colour-preserving” is replaced with “colour-permuting.” This is reminiscent of the CI (Cayley Isomorphism) property [8], and this similarity motivated our choice of terminology.

We thank Gabriel Verret for pointing out that the quaternion group Q_8 is not CCA. In fact, two different groups of order 8 are not CCA:

Example 2.2 (G. Verret). $\mathbb{Z}_4 \times \mathbb{Z}_2$ and Q_8 are not CCA.

Proof. (Q_8) Let $\Gamma = \text{Cay}(Q_8; \{\pm i, \pm j\})$. This is the complete bipartite graph $K_{4,4}$. (See Figure 1 with the labels that are inside the vertices.) Let φ be the graph automorphism that interchanges the vertices k and $-k$ while fixing every other vertex. This is clearly not an automorphism of G since i and j are fixed by φ and generate G , but $\varphi \neq 1$. It is, however, a colour-preserving automorphism of Γ .

($\mathbb{Z}_4 \times \mathbb{Z}_2$) Let $\Gamma = \text{Cay}(\mathbb{Z}_4 \times \mathbb{Z}_2; \{\pm(1, 0), \pm(1, 1)\})$. This is again the complete bipartite graph $K_{4,4}$. (See Figure 1 with the labels that are outside the vertices.) Let φ be the graph automorphism that interchanges the vertices $(0, 1)$ and $(2, 1)$ while fixing all of the other vertices. This is clearly not an automorphism of G since $(1, 0)$ and $(1, 1)$ are fixed by φ and generate G , but $\varphi \neq 1$. It is, however, a colour-preserving automorphism of Γ . $\square$

Both of the groups in Example 2.2 are generalized dicyclic (cf. Definition 2.6):

- Q_8 is the generalized dicyclic group over $\mathbb{Z}_4$, and
- $\mathbb{Z}_4 \times \mathbb{Z}_2$ is the generalized dicyclic group over $\mathbb{Z}_2 \times \mathbb{Z}_2$.

More generally, we will see in Corollary 2.8(2) below that no generalized dicyclic group is CCA, except $\mathbb{Z}_4$.

We will see in Theorem 6.8 that the following example is the smallest group of odd order that is not CCA.




Figure 1: Interchanging the two black vertices while fixing all of the white vertices is a colour-preserving graph automorphism that fixes the identity vertex but is not a group automorphism.

Example 2.3. The nonabelian group of order 21 is not CCA.

Proof. Let $G = \langle a, x \mid a^3 = e, a^{-1}xa = x^2 \rangle$. (Since $x = e^{-1}xe = a^{-3}xa^3 = x^8$, the relations imply $x^7 = e$, so G has order 21.) By letting $b = ax$, we see that G also has the presentation

$$G = \langle a, b \mid a^3 = e, (ab^{-1})^2 = b^{-1}a \rangle.$$

As illustrated in Figure 2, every element of G can be written uniquely in the form

$$a^i b^j a^k, \text{ where } i, j, k \in \{0, \pm 1\} \text{ and } j = 0 \Rightarrow k = 0.$$

Define

$$\varphi(a^i b^j a^k) = \begin{cases} b^j a^{-k} & \text{if } i = 0, \\ ab^{-j} a^k & \text{if } i = 1, \\ a^{-1} b^{-j} a^{-k} & \text{if } i = -1. \end{cases}$$

Then φ is a colour-preserving automorphism of $\text{Cay}(G; \{a^{\pm 1}, b^{\pm 1}\})$ (see Figure 2). However, φ is not affine, since it fixes e , but is not an automorphism of G (because $\varphi(ab) = ab^{-1} \neq ab = \varphi(a)\varphi(b)$). $\square$

See Proposition 3.3 for a generalization of the following example.

Example 2.4. The wreath product $\mathbb{Z}_m \wr \mathbb{Z}_n$ is not CCA whenever $m \geq 3$ and $n \geq 2$.

Proof. This group is a semidirect product

$$(\mathbb{Z}_m \times \mathbb{Z}_m \times \cdots \times \mathbb{Z}_m) \rtimes \mathbb{Z}_n.$$

For the generators $a = ((1, 0, 0, \dots, 0), 0)$ and $b = ((0, 0, \dots, 0), 1)$, the map

$$((x_1, x_2, x_3, \dots, x_n), y) \mapsto ((-x_1, x_2, x_3, \dots, x_n), y)$$




Figure 2: The colour-preserving automorphism φ fixes every black vertex, but interchanges the two vertices labeled $\textcircled{i}$, for $1 \leq i \leq 8$. Since the neighbours of both copies of $\textcircled{i}$ have the same labels (for example, the vertices labeled $\textcircled{7}$ are connected by a black edge to $\textcircled{1}$ and $\textcircled{5}$, and by a white edge to $\textcircled{6}$ and $\textcircled{8}$), we see that φ is indeed a colour-preserving automorphism of the graph (if the orientations of the edges are ignored).

(negate a single factor of the abelian normal subgroup) is a colour-preserving automorphism of $\text{Cay}(\mathbb{Z}_m \wr \mathbb{Z}_n; \{a^{\pm 1}, b^{\pm 1}\})$ that fixes the identity element but is not a group automorphism. $\square$

The following construction provides many additional examples of non-CCA groups by generalizing the idea of Example 2.2.

Proposition 2.5. Suppose there is a generating set S of G , an element τ of G , and a subset T of S , such that:

- $S = S^{-1}$,
- τ is an element of order 2,
- each element of S is either centralized or inverted by τ ,
- $t^2 = \tau$ for all $t \in T$,
- the subgroup $\langle (S \setminus T) \cup \{\tau\} \rangle$ is not all of G , and
- either $|G : \langle (S \setminus T) \cup \{\tau\} \rangle| > 2$ or τ is not in the centre of G .

Then G is not CCA.

Proof. For convenience, let $H = \langle (S \setminus T) \cup \{\tau\} \rangle$. Since $\langle S \rangle = G$, but, by assumption, $H \neq G$, there exists some $x \in T \setminus H$. Define

$$\varphi(g) = \begin{cases} g\tau & \text{if } g \in xH, \\ g & \text{otherwise.} \end{cases}$$

It is obvious that φ fixes e , since $e \notin xH$.

We claim that φ is not an automorphism of G . If $|G : H| > 2$, this follows from the fact that a nonidentity automorphism cannot fix more than half of the elements of G . Thus, we may assume $|G : H| = 2$. Then, by assumption, there is some element h of G that does not commute with τ . Since τ commutes with every element of T (because $\tau = t^2$), we see that we may assume $h \in H$. If φ is an automorphism, then, since it is the identity on the normal subgroup H of G , but $x^{-1} = xx^{-2} = x\tau \in xH$, we have:

$$x^{-1}hx = \varphi(x^{-1}hx) = \varphi(x^{-1}) \cdot \varphi(h) \cdot \varphi(x) = x^{-1}\tau \cdot h \cdot x\tau \neq x^{-1}hx\tau^2 = x^{-1}hx.$$

This is a contradiction.

Since each element of S is either centralized or inverted by τ , we know that right-multiplication by τ is a colour-preserving automorphism of $\text{Cay}(G; S)$. Restricting to xH , this tells us that φ preserves colours (and existence) of all edges of $\text{Cay}(G; S)$ that have both endvertices in xH .

Now consider an edge from g to h , where $g \in xH$ and $h \notin xH$. There is some element $t \in T$ such that $gt = h$, and there is an edge of the same colour from $\varphi(g) = g\tau$ to $g\tau t^{-1}$. Since $t^2 = \tau$ and $\tau^2 = e$, we have $t^{-1} = \tau t$. Hence, the edge is from $\varphi(g)$ to

$$g\tau t^{-1} = gt^2 t^{-1} = gt = h = \varphi(h).$$

Thus φ preserves the existence and colour of every edge from a vertex in xH to a vertex outside of xH . Since the only vertices moved by φ are in xH , this shows that φ is a colour-preserving automorphism of $\text{Cay}(G; S)$. $\square$

Definition 2.6. Let A be an abelian group of even order. Choose an involution y of A . The corresponding *generalized dicyclic group* is

$$\text{Dic}(A, y) = \langle x, A \mid x^2 = y, x^{-1}ax = a^{-1}, \forall a \in A \rangle.$$

Definition 2.7. For $n \geq 1$, let

$$\text{SemiD}_{16n} = \langle a, x \mid a^{8n} = x^2 = e, xa = a^{4n-1}x \rangle.$$

This is a *semidihedral* (or *quasidihedral*) group. The term is usually used only when n is a power of 2, but the construction is valid more generally.

We have already seen in Example 2.2 that $\mathbb{Z}_4 \times \mathbb{Z}_2$ and Q_8 are not CCA. Here are a few additional examples that come from Proposition 2.5.

Corollary 2.8. The following groups are not CCA:

1. $\mathbb{Z}_{2^k} \times \mathbb{Z}_2 \times \mathbb{Z}_2$, for any $k \geq 2$,
2. every generalized dicyclic group except $\mathbb{Z}_4$, and

3. every semidihedral group.

Proof. (1) Apply Proposition 2.5 with $\tau = (2^{k-1}, 0, 0)$, $T = \{(2^{k-2}, 1, 0), (2^{k-2}, 0, 1)\}$, and $S = \{(1, 0, 0)\} \cup T$.

(2) For $G = \text{Dic}(A, y) = \langle x, y, A \rangle$, apply Proposition 2.5 with $\tau = y$ and $S = T = xA$. (We have $|G : \langle (S \setminus T) \cup \{\tau\} \rangle| = |G : \langle \tau \rangle| = |G|/2 > 2$, since $G \not\cong \mathbb{Z}_4$.)

(3) For $G = \text{SemiD}_{16n} = \langle a, x \rangle$, apply Proposition 2.5 with $\tau = a^{4n}$, $T = \{(ax)^{\pm 1}\}$, and $S = \{x\} \cup T$. (Note that $|G : \langle (S \setminus T) \cup \{\tau\} \rangle| = |G : \langle x, \tau \rangle| = |G|/4 \geq 4$.) $\square$

3 Direct products and semidirect products

Proposition 3.1. If G_1 is not strongly CCA, and G_2 is any group, then $G_1 \times G_2$ is not strongly CCA. Furthermore, the same is true with “CCA” in the place of “strongly CCA.”

Proof. Since G_1 is not strongly CCA, some connected Cayley graph $\text{Cay}(G_1; S_1)$ on G_1 has a colour-permuting automorphism φ_1 that is not affine. Let π be a permutation of S_1 , such that $\varphi_1(g_1 s) \in \{\varphi_1(g_1) \pi(s)^{\pm 1}\}$ for all $g_1 \in G_1$ and $s \in S_1$. (If G_1 is not CCA, then we may assume π is the identity permutation.) Now, fix any connected Cayley graph $\text{Cay}(G_2; S_2)$ on G_2 , and let

$$S = (S_1 \times \{e\}) \cup (\{e\} \times S_2),$$

so $\text{Cay}(G_1 \times G_2; S)$ is connected. (It is isomorphic to the Cartesian product $\text{Cay}(G_1; S_1) \square \text{Cay}(G_2; S_2)$.)

Define a permutation φ of $G_1 \times G_2$ by $\varphi(x_1, x_2) = (\varphi_1(x_1), x_2)$. For all $(x_1, x_2) \in G_1 \times G_2$ and $s_i \in S_i$, we have

- $\varphi((x_1, x_2) \cdot (s_1, e)) = (\varphi_1(x_1 s_1), x_2) \in \{\varphi(x_1, x_2) \cdot (\pi(s_1), e)^{\pm 1}\}$, and
- $\varphi((x_1, x_2) \cdot (e, s_2)) = (\varphi_1(x_1), x_2 s_2) = \varphi(x_1, x_2) \cdot (e, s_2)$.

Therefore, φ is a colour-permuting automorphism of $\text{Cay}(G_1 \times G_2; S)$ (and it is colour-preserving if π is the identity permutation of S_1).

To complete the proof that $G_1 \times G_2$ is not strongly CCA (and is not CCA if π is the identity permutation of S_1), it suffices to show that φ is not affine. We prove this by contradiction: suppose there exists an automorphism α of $G_1 \times G_2$ and $(g_1, g_2) \in G_1 \times G_2$, such that $\varphi(x_1, x_2) = \alpha(g_1 x_1, g_2 x_2)$, for all $(x_1, x_2) \in G_1 \times G_2$. By the definition of φ , this implies $\varphi_1(x_1) = \alpha(g_1 x_1)$ for all $x_1 \in G_1$. Since $\varphi_1(x_1) \in G_1$ and $g_1 x_1$ is an arbitrary element of G_1 , we conclude that $\alpha(G_1) \subseteq G_1$, so the restriction of α to G_1 is an automorphism of G_1 . Hence, the equation $\varphi_1(x_1) = \alpha(g_1 x_1)$ implies that φ_1 is affine. This contradicts the choice of φ_1 . $\square$

Proposition 3.1 tells us that if $G_1 \times G_2$ is CCA, then G_1 and G_2 must both be CCA. The converse is not true. (For example, $\mathbb{Z}_4$ and $\mathbb{Z}_2$ are both CCA, but Example 2.2 tells us that the direct product $\mathbb{Z}_4 \times \mathbb{Z}_2$ is not CCA.) However, the converse is indeed true when the groups are of relatively prime order:

Proposition 3.2. Assume $\gcd(|G_1|, |G_2|) = 1$. Then $G_1 \times G_2$ is CCA (or strongly CCA) if and only if G_1 and G_2 are both CCA (or strongly CCA, respectively).

Proof. ($\Rightarrow$) Proposition 3.1.

($\Leftarrow$) Let

- $G = G_1 \times G_2$,
- S be a generating set of G ,
- φ be a colour-permuting automorphism of $\text{Cay}(G; S)$ that fixes the identity element (see Remark 2.1),
- $\pi_i: G_1 \times G_2 \rightarrow G_i$ be the natural projection, and
- k be a multiple of $|G_2|$ that is $\equiv 1 \pmod{|G_1|}$, so $g^k = \pi_1(g)$ for all $g \in G$.

Consider some $s \in S$, and let $t = \varphi(s)$, so $\varphi(xs^i) = \varphi(x)t^{\pm i}$ for all $x \in G$ and $i \in \mathbb{Z}$. Then, for all $g \in G$, we have

$$\varphi(g \pi_1(s)) = \varphi(gs^k) = \varphi(g)t^{\pm k} = \varphi(g) \cdot \pi_1(t)^{\pm 1}. \quad (*)$$

Since $\pi_1(S)$ generates G_1 , this implies there is a well-defined permutation φ_2 of G_2 , such that

$$\varphi(G_1 \times \{g_2\}) = G_1 \times \{\varphi_2(g_2)\} \text{ for all } g_2 \in G_2.$$

By repeating the argument with the roles of G_1 and G_2 interchanged, we conclude that there is a permutation φ_1 of G_1 , such that

$$\varphi(g_1, g_2) = (\varphi_1(g_1), \varphi_2(g_2)) \text{ for all } (g_1, g_2) \in G_1 \times G_2.$$

Now, (*) implies that φ_1 is a colour-permuting automorphism of $\text{Cay}(G_1; \pi_1(S))$. Similarly, φ_2 is a colour-permuting automorphism of $\text{Cay}(G_2; \pi_2(S))$. Since each G_i is CCA, we conclude that φ_i is an automorphism of G_i . So φ is an automorphism of $G_1 \times G_2$. $\square$

The idea used in Example 2.4 yields the following result that generalizes the CCA part of Proposition 3.1.

Proposition 3.3. Suppose $G = H \rtimes K$ is a semidirect product, and $\text{Cay}(H; S_0)$ is a connected Cayley graph of H , such that:

- S_0 is invariant under conjugation by every element of K , and
- there is a colour-preserving automorphism φ_0 of $\text{Cay}(H; S_0)$, such that either
 - φ_0 is not affine, or
 - $\varphi_0(e) = e$, and there exist $s \in S_0$ and $k \in K$, such that $\varphi_0(k^{-1}sk) \neq k^{-1}\varphi_0(s)k$.

Then G is not CCA.

Proof. Define $\varphi: G \rightarrow G$ by $\varphi(hk) = \varphi_0(h)k$. We claim that φ is a colour-preserving automorphism of $\text{Cay}(G; S_0 \cup K)$ that is not affine (so G is not CCA, as desired).

For $h \in H$ and $k, k_1 \in K$, we have

$$\varphi(hk k_1) = \varphi_0(h) k k_1 = \varphi(hk) k_1,$$

so φ preserves the colour of K -edges. Now consider some $s \in S_0$ and let ${}^k s = k s k^{-1} \in S_0$. Then, since φ_0 is colour preserving, we have

$$\varphi(hk s) = \varphi(h {}^k s k) = \varphi_0(h {}^k s) k = (\varphi_0(h) ({}^k s)^{\pm 1}) k = \varphi_0(h) k s^{\pm 1} = \varphi(hk) s^{\pm 1},$$

so φ also preserves the colour of S_0 -edges. Hence, φ is colour-preserving.

Now, suppose φ is affine. Then the restriction φ_0 of φ to H is also affine, so, by assumption, we must have $\varphi(e) = e$, so φ is an automorphism of G . Hence, for all $s \in S_0$ and $k \in K$, we have

$$\varphi_0(k^{-1} s k) = \varphi(k^{-1} s k) = \varphi(k)^{-1} \varphi(s) \varphi(k) = k^{-1} \varphi(s) k = k^{-1} \varphi_0(s) k.$$

This contradicts the hypotheses of the proposition. $\square$

Remark 3.4. Proposition 3.3 can be generalized slightly: assume $G = HK$ and $H \triangleleft G$ (but do not assume $H \cap K = \{e\}$, which would make G a semidirect product). Then the above proof applies if we make the additional assumption that $\varphi_0(hk) = \varphi_0(h) k$ for all $h \in H$ and $k \in H \cap K$.

4 Abelian groups

The following result shows that all non-CCA abelian groups can be constructed from examples that we have already seen in Example 2.2 and Corollary 2.8 (and that CCA and strongly CCA are equivalent for abelian groups).

Proposition 4.1. For an abelian group G , the following are equivalent:

1. G has a direct factor that is isomorphic to either $\mathbb{Z}_4 \times \mathbb{Z}_2$ or a group of the form $\mathbb{Z}_{2^k} \times \mathbb{Z}_2 \times \mathbb{Z}_2$, with $k \geq 3$.
2. G is not CCA.
3. G is not strongly CCA.

Proof. (1 $\Rightarrow$ 2) This is immediate from Example 2.2, Corollary 2.8, and Proposition 3.1.

(2 $\Rightarrow$ 3) Obvious.

(3 $\Rightarrow$ 1) We prove the contrapositive. Assume G does not have any direct summands of the form specified in (1). Given a connected Cayley graph $\text{Cay}(G; S)$ on G , and a colour-permuting automorphism φ of $\text{Cay}(G; S)$, such that $\varphi(0) = 0$, we will show that φ is an automorphism of G .

From Proposition 3.2 (and the fact that every abelian group is the direct sum of its Sylow subgroups), we may assume G is a p -group for some prime p . Then

$$G \cong \mathbb{Z}_{p^{k_1}} \times \mathbb{Z}_{p^{k_2}} \times \cdots \times \mathbb{Z}_{p^{k_m}}, \quad \text{with } k_1 \geq k_2 \geq \cdots \geq k_m \geq 1.$$

Since S is a generating set, it is easy to see that there is some $s_1 \in S$, such that $|s_1| = p^{k_1}$. Also, it is a basic fact about finite abelian groups that every cyclic subgroup of maximal order is a direct summand [4, Lem. 1.3.3, p. 10]. Therefore, by induction on i , we see that there exist $s_1, \dots, s_m \in S$, such that if we let $G_i = \langle s_1, \dots, s_i \rangle$, then

$$G_i \cong G_{i-1} \times \mathbb{Z}_{p^{k_i}} \quad \text{and} \quad G \cong G_i \times \mathbb{Z}_{p^{k_{i+1}}} \times \cdots \times \mathbb{Z}_{p^{k_m}}, \quad \text{for each } i.$$

It is important to note that each element of G_i can be written uniquely in the form

$$g + rs_i, \text{ with } g \in G_{i-1} \text{ and } -p^{k_i}/2 < r \leq p^{k_i}/2 \text{ (and } r \in \mathbb{Z}). \quad (\dagger)$$

For convenience, also let

$$t_i = \varphi(s_i) \quad \text{and} \quad H_i = \langle t_1, \dots, t_i \rangle.$$

We will show, by induction on i , that H_i is a direct factor of G , and the restriction of φ to G_i is an isomorphism onto H_i . (Note that this implies $G/G_i \cong G/H_i$, by the uniqueness of the decomposition of G as a direct sum of cyclic groups.) Taking $i = m$ yields the desired conclusion that φ is an automorphism of G .

The base case $i = 0$ is trivial. For the induction step, write $G = G_{i-1} \times \overline{G}$, so

$$\overline{G} \cong G/G_{i-1} \cong \mathbb{Z}_{p^{k_i}} \times \mathbb{Z}_{p^{k_{i+1}}} \times \dots \times \mathbb{Z}_{p^{k_m}},$$

and let $\bar{\cdot}: G \rightarrow \overline{G}$ be the natural projection. Then $\langle \overline{s_i} \rangle = \overline{G_i} \cong \mathbb{Z}_{p^{k_i}}$ is a direct summand of $\overline{G}$. Since φ is colour-permuting (and $H_{i-1} = \varphi(G_{i-1})$ is a subgroup), it is easy to see that the order of t_i in G/H_{i-1} is equal to p^{k_i} (the same as the order of s_i in G/G_{i-1}), and that $\varphi(p^{k_i}s_i) = p^{k_i}t_i$. This implies that if we define

$$\alpha: G_i \rightarrow H_i \text{ by } \alpha(g + rs_i) = \varphi(g) + rt_i \quad \text{for } g \in G_{i-1} \text{ and } r \in \mathbb{Z},$$

then α is a well-defined isomorphism. So we need only show that the restriction of φ to G_i is equal to α .

Suppose $\varphi|_{G_i} \neq \alpha$. (This will lead to a contradiction.) Since φ is colour-permuting and, by definition, α agrees with φ on G_{i-1} , this implies there is some $g \in G_{i-1}$, such that $\varphi(g + s_i) \neq \alpha(g + s_i)$. However, since φ is colour-permuting, we know

$$\varphi(g + s_i) = \varphi(g) \pm \varphi(s_i) = \alpha(g) \pm t_i.$$

Since $\alpha(g + s_i) = \alpha(g) + t_i$, the preceding two sentences imply

$$\varphi(g + s_i) = \alpha(g) - t_i \in H_{i-1} - t_i.$$

Furthermore, since φ is colour-permuting (and $\varphi(s_j) = t_j$), we know that it maps edges of colour $\{s_1^{\pm 1}\}, \dots, \{s_{i-1}^{\pm 1}\}$ to edges of colour $\{t_1^{\pm 1}\}, \dots, \{t_{i-1}^{\pm 1}\}$, so

$$\varphi(x + h) \in \varphi(x) + H_{i-1} \text{ for all } x \in G \text{ and } h \in H_{i-1}.$$

Taking $x = s_i$ and $h = g$ yields

$$\varphi(g + s_i) \in H_{i-1} + \varphi(s_i) = H_{i-1} + t_i.$$

This contradicts the uniqueness of r in the analogue of $(\dagger)$ for H_i , unless $1 = p^{k_i}/2$. Hence, we must have $p^{k_i} = 2$ (so $\mathbb{Z}_2$ is a direct summand of G), which means $p = 2$ and $k_i = 1$.

We have

$$\begin{aligned} \varphi(g) + 2t_i &= \alpha(g + 2s_i) && \text{(definition of } \alpha) \\ &= \varphi(g + 2s_i) && (g + 2s_i = g + p^{k_i}s_i \in G_{i-1}) \\ &= \varphi(g) - 2t_i && (\varphi(g + s_i) = \alpha(g) - t_i = \varphi(g) - t_i), \end{aligned}$$

so $4t_i = 0$. Also note that, since

$$\varphi(g) + t_i = \alpha(g + s_i) \neq \varphi(g + s_i) = \varphi(g) - t_i,$$

we must have $2t_i \neq 0$. So $|t_i| = 4$.

Since $\langle s_1, \dots, s_{i-1} \rangle = G_{i-1}$, there must exist $g' \in G_{i-1}$, and $j < i$, such that

$$\varphi(g' + s_i) = \alpha(g') + t_i, \text{ but } \varphi(g' + s_j + s_i) = \alpha(g' + s_j) - t_i = \alpha(g') + t_j - t_i.$$

Since φ is colour-permuting, we also have

$$\varphi(g' + s_j + s_i) = \varphi(g' + s_i) \pm t_j = \alpha(g') + t_i \pm t_j.$$

Hence, $t_j - t_i = t_i \pm t_j$, so $t_j \mp t_j = 2t_i$. Since $2t_i \neq 0$, we conclude that $2t_j = 2t_i$; hence, $|t_j| = 4$.

Since $2^{k_j} = |H_j : H_{j-1}|$ is a divisor of $|t_j|$, and $|t_j| = 4$, there are two possibilities for k_j :

- If $k_j = 2$, then $\mathbb{Z}_4 \times \mathbb{Z}_2 \cong \mathbb{Z}_{2^{k_j}} \times \mathbb{Z}_{2^{k_i}}$ is a direct summand of G .
- If $k_j = 1$, then, since $|t_j| = 4$, there must be some $\ell < j$, such that $k_\ell \geq 2$. This implies that $\mathbb{Z}_{2^{k_\ell}} \times \mathbb{Z}_2 \times \mathbb{Z}_2 \cong \mathbb{Z}_{2^{k_\ell}} \times \mathbb{Z}_{2^{k_j}} \times \mathbb{Z}_{2^{k_i}}$ is a direct summand of G .

Each of these possibilities contradicts our assumption that there are no direct summands as specified in (1) of the statement of the proposition. $\square$

Corollary 4.2. For $n \in \mathbb{Z}^+$, there is a non-CCA abelian group of order n if and only if n is divisible by 8.

5 Generalized dihedral groups

Definition 5.1. The *generalized dihedral group* over an abelian group A is the group

$$\langle \sigma, A \mid \sigma^2 = e, \sigma a \sigma = a^{-1}, \forall a \in A \rangle.$$

Lemma 5.2. Suppose D is the generalized dihedral group over an abelian group A , and φ is a colour-permuting automorphism of a connected Cayley graph $\text{Cay}(D; S)$, such that $\varphi(e) = e$. If A is strongly CCA, and $\varphi(S \cap A) = S \cap A$, then φ is an automorphism of D .

Proof. Label the elements of S as $S = \{a_1, a_2, \dots, a_k, \sigma_1, \sigma_2, \dots, \sigma_t\}$, where $a_i \in A$ for $1 \leq i \leq k$, and $\sigma_i \notin A$ for $1 \leq i \leq t$ (so each σ_i is an involution whose action by conjugation inverts every element of A). By assumption, $\{a_1, a_2, \dots, a_k\}$ and $\{\sigma_1, \sigma_2, \dots, \sigma_t\}$ are invariant under φ . Thus, for each i , we have

- $\varphi(a_i) = a'_i$ for some $a'_i \in \{a_1, a_2, \dots, a_k\}$, and
- $\varphi(\sigma_i) = \sigma'_i$ for some $\sigma'_i \in \{\sigma_1, \sigma_2, \dots, \sigma_t\}$.

Notice that since $\sigma_1, \dots, \sigma_t$ are involutions, each σ_i is its own inverse. Therefore, whenever σ is a word in $\sigma_1, \dots, \sigma_t$ and $g \in D$, the fact that φ is a colour-permuting automorphism means that $\varphi(g\sigma) = \varphi(g)\sigma'$, where σ' is formed from σ by replacing each instance of σ_i in σ by σ'_i . Therefore, if we let Σ be the subgroup generated by $\{\sigma_1, \dots, \sigma_t\}$,

then φ is a colour-preserving automorphism of the Cayley graph $\text{Cay}(D; S \cup \Sigma)$. Hence, there is no harm in assuming that $S = S \cup \Sigma$, so $\Sigma \subseteq S$.

Since $\langle S \cap A \rangle$ is normal in D (in fact, every subgroup of A is normal, because every element of D either centralizes or inverts it), we have $D = \langle S \cap A \rangle \Sigma$. Therefore $A = \langle S \cap A \rangle (\Sigma \cap A) = \langle S \cap A \rangle$, so $\text{Cay}(A; S \cap A)$ is connected. Since φ is colour-preserving, and $\varphi(S \cap A) = S \cap A$, this implies that $\varphi(A) = A$. So φ is a colour-permuting automorphism of the connected Cayley graph $\text{Cay}(A; S \cap A)$. Since, by assumption, A is strongly CCA, this implies that $\varphi|_A$ is an automorphism of A . So $\varphi(ab^\epsilon) = \varphi(a) \varphi(b)^\epsilon$ for all $a, b \in A$ and $\epsilon \in \mathbb{Z}$.

Now we are ready to show that φ is an automorphism of D . Let $g, h \in D$. Then we may write $g = a\sigma$ and $h = b\tilde{\sigma}$, where $a, b \in A$ and $\sigma, \tilde{\sigma} \in \{e, \sigma_1\}$. For convenience, let $\epsilon \in \{\pm 1\}$, such that $\sigma c \sigma = c^\epsilon$ for all $c \in A$. Note that, since $\sigma'_1 \in \{\sigma_1, \dots, \sigma_t\}$, we know that σ_1 and σ'_1 both invert A , so we also have $\sigma' c \sigma' = c^\epsilon$. Then

$$\varphi(gh) = \varphi(a\sigma \cdot b\tilde{\sigma}) = \varphi(ab^\epsilon \cdot \sigma\tilde{\sigma}) = \varphi(a) \varphi(b)^\epsilon \cdot \sigma'\tilde{\sigma}' = \varphi(a)\sigma' \cdot \varphi(b)\tilde{\sigma}' = \varphi(g) \cdot \varphi(h).$$

Since $g, h \in D$ are arbitrary, this proves that φ is an automorphism of D . $\square$

Proposition 5.3. The generalized dihedral group D over an abelian group A is CCA if and only if A is CCA.

Proof. ($\Leftarrow$) Note that if φ is any colour-preserving automorphism of a connected Cayley graph $\text{Cay}(D; S)$ such that $\varphi(e) = e$, then $\varphi(S \cap A) = S \cap A$, since A is closed under inverses. Furthermore, A is strongly CCA, since it is assumed to be CCA and every CCA abelian group is strongly CCA (see Proposition 4.1). Therefore, Lemma 5.2 implies that φ is a group automorphism. So D is CCA.

($\Rightarrow$) Write $D = A \rtimes \langle \sigma \rangle$. Since A is not CCA, there is a colour-preserving automorphism φ_0 of some connected Cayley graph $\text{Cay}(A; S)$, such that φ_0 is not affine. Since σ inverts every element of S , it is easy to see that $\text{Cay}(D; S \cup \{\sigma\})$ is isomorphic to the Cartesian product $\text{Cay}(A; S) \square P_2$. So the proof of Proposition 3.1 provides a colour-preserving automorphism φ of $\text{Cay}(D; S \cup \{\sigma\})$ whose restriction to A is φ_0 , which is not an affine map. Therefore, φ is not affine. $\square$

The following result is the special case where A is cyclic (since Proposition 4.1 implies that every cyclic group is CCA).

Corollary 5.4. Every dihedral group is CCA.

Lemma 5.5. If T is a generating set of a group H , and σ is a nontrivial automorphism of H , such that $\sigma(t) \in \{t^{\pm 1}\}$ for every $t \in T$, then the group $G = (H \rtimes \langle \sigma \rangle) \times \mathbb{Z}_2$ is not strongly CCA.

Proof. Let $G' = H \times \mathbb{Z}_2 \times \mathbb{Z}_2$ and define $\varphi: G \rightarrow G'$ by $\varphi(h, \sigma^x, y) = (h, x, y)$ for $h \in H$ and $x, y \in \mathbb{Z}_2$. Since $\sigma(t) \in \{t^{\pm 1}\}$ for every t , it is easy to verify that φ is a colour-respecting isomorphism

$$\begin{aligned} & \text{from } \text{Cay}(G; (H, e, 0) \cup \{(e, \sigma, 0), (e, 0, 1)\}) \\ & \text{to } \text{Cay}(G'; (H, 0, 0) \cup \{(e, 1, 0), (e, 0, 1)\}). \end{aligned}$$

Permuting the two $\mathbb{Z}_2$ factors of G' provides an automorphism of G' that preserves the generating set, and therefore corresponds to a colour-permuting automorphism of the two

Cayley graphs. However, it is not an automorphism of G , since it takes the central element $(e, e, 1)$ to $(e, \sigma, 0)$, which is not central (since the automorphism σ is nontrivial). $\square$

Proposition 5.6. The generalized dihedral group over an abelian group A is strongly CCA if and only if either A does not have $\mathbb{Z}_2$ as a direct factor, or A is an elementary abelian 2-group (in which case, the generalized dihedral group is also an elementary abelian 2-group).

Proof. ($\Rightarrow$) Suppose $A = A' \times \mathbb{Z}_2$, and A' is not elementary abelian. Then the generalized dihedral group $A \rtimes \langle \sigma \rangle$ over A is isomorphic to $(A' \rtimes \langle \sigma \rangle) \times \mathbb{Z}_2$, so Lemma 5.5 tells us that it is not strongly CCA.

($\Leftarrow$) Let $D = A \rtimes \langle \sigma \rangle$ be the generalized dihedral group over A , and let φ be a colour-permuting automorphism of a connected Cayley graph $\text{Cay}(D; S)$, such that $\varphi(e) = e$. We may assume A does not have $\mathbb{Z}_2$ as a direct factor (otherwise, the desired conclusion follows from the fact that every elementary abelian 2-group is strongly CCA (see Proposition 4.1)). From Proposition 4.1, we see that A is strongly CCA. Hence, the desired conclusion will follow from Lemma 5.2 if we show that $\varphi(S \cap A) = S \cap A$.

Let $a \in S \cap A$. Since φ is colour-permuting, we have $|\varphi(s)| = |s|$ for all $s \in S$. Also, we know that $|g| = 2$ for all $g \in D \setminus A$. Therefore, it is obvious that $\varphi(a) \in S \cap A$ if $|a| \neq 2$.

So we may assume $|a| = 2$. Since A does not have $\mathbb{Z}_2$ as a direct factor, this implies that a is a square in A : that is, we have $a = x^2$, for some $x \in A$. Also, since $\text{Cay}(D; S)$ is connected, we may write $x = s_1 s_2 \cdots s_n$ for some $s_1, \dots, s_n \in S$. So $a = (s_1 s_2 \cdots s_n)^2$ can be written as a word in which every element of S occurs an even number of times. Since φ is colour-permuting, this implies that $\varphi(a)$ can be written as a word in which, for each $s \in S$, the total number of occurrences of either s or s^{-1} is even. Since s and s^{-1} both either centralize A or invert it, this implies that $\varphi(a)$ centralizes A . Since A is self-centralizing in D , we conclude that $\varphi(a) \in A$, as desired. $\square$

6 Groups of odd order

The following notation will be assumed throughout this section.

Notation 6.1. For a fixed Cayley graph $\text{Cay}(G; S)$:

- $\mathcal{A}^0$ is the group of all colour-preserving automorphisms of $\text{Cay}(G; S)$.
- $\widehat{G}$ is the subgroup of $\mathcal{A}^0$ consisting of all left translations by elements of G . (Although we do not need this terminology, it is often called the *left regular representation* of G .)
- H_e is the stabilizer of the identity element e in $\text{Cay}(G; S)$, for any subgroup H of $\mathcal{A}^0$.

Remark 6.2. It is well known (and very easy to prove) that a permutation of G is affine if and only if it normalizes $\widehat{G}$ (see, for example [11, Lem. 2]).

Lemma 6.3. $\mathcal{A}_e^0$ is a 2-group.

Proof. Let $\varphi \in \mathcal{A}_e^0$, so φ is a colour-preserving automorphism of $\text{Cay}(G; S)$ that fixes e . If C is any monochromatic cycle through e , then either φ is the identity on C or φ reverses

the orientation of C . Therefore, φ^2 acts trivially on the union of all monochromatic cycles that contain e . This implies that φ^2 acts trivially on all vertices at distance ≤ 1 from e .

Repeating the argument shows that φ^{2^k} acts trivially on all vertices at distance $\leq k - 1$ from e . For k larger than the diameter of $\text{Cay}(G; S)$, this implies that φ^{2^k} is trivial. So the order of φ is a power of 2. $\square$

Proposition 6.4. Let $\text{Cay}(G; S)$ be a connected Cayley graph on a group G of odd order. If $\text{Cay}(G; S)$ is CCA, then $\text{Cay}(G; S)$ is strongly CCA.

Proof. Let $\mathcal{A}^\bullet$ be the group of all colour-permuting automorphisms of $\text{Cay}(G; S)$. Since $\mathcal{A}^\bullet$ acts on the set of colours, and $\mathcal{A}^0$ is the kernel of this action (and the kernel of a homomorphism is always normal), it is obvious that $\mathcal{A}^0 \triangleleft \mathcal{A}^\bullet$. Also, since $\text{Cay}(G; S)$ is CCA, we have $\widehat{G} \triangleleft \mathcal{A}^0$ (cf. Remark 6.2). Furthermore, $|G|$ is odd, $|\mathcal{A}_e^0|$ is a power of 2, and $\mathcal{A}^0 = \widehat{G} \cdot \mathcal{A}_e^0$. Therefore, $\widehat{G}$ is the (unique) largest normal subgroup of odd order in $\mathcal{A}^0$. The uniqueness implies that $\widehat{G}$ is *characteristic* in $\mathcal{A}^0$. (That is, it is fixed by all automorphisms of $\mathcal{A}^0$.) So $\widehat{G}$ is a characteristic subgroup of the normal subgroup $\mathcal{A}^0$ of $\mathcal{A}^\bullet$. Since every characteristic subgroup of a normal subgroup is normal [4, Thm. 2.1.2(ii), p. 16], this implies $\widehat{G} \triangleleft \mathcal{A}^\bullet$. Therefore G is strongly CCA (see Remark 6.2). $\square$

Wreath products $\mathbb{Z}_m \wr \mathbb{Z}_n$ provide examples of non-CCA groups of odd order (see Example 2.4). We will see in Theorem 6.8 that the following slightly more general construction is essential for understanding many of the other non-CCA groups of odd order.

Example 6.5. Let α be an automorphism of a group A , and let $n \in \mathbb{Z}^+$. Then we can define an automorphism $\tilde{\alpha}$ of A^n by

$$\tilde{\alpha}(w_1, \dots, w_n) = (\alpha(w_n), w_1, w_2, \dots, w_{n-1}).$$

It is easy to see that the order of $\tilde{\alpha}$ is n times the order of α , so we may form the corresponding semidirect product $A^n \rtimes \mathbb{Z}_{n|\alpha|}$. Let us call this the *semi-wreathed product* of A by $\mathbb{Z}_n$, with respect to the automorphism α , and denote it $A \wr_\alpha \mathbb{Z}_n$. (If α is the trivial automorphism, then this is the usual wreath product $A \wr \mathbb{Z}_n$.)

Negating the first coordinate, as in Example 2.4, shows that if $n > 1$ and A is abelian, but not an elementary abelian 2-group, then $A \wr_\alpha \mathbb{Z}_n$ is not CCA.

Remark 6.6. Because it may be of interest to find minimal examples, we point out that every semi-wreathed product of odd order satisfying the conditions in the final paragraph of Example 6.5 must contain a subgroup that is isomorphic to a semi-wreathed product $A \wr_\alpha \mathbb{Z}_q$, where A is an elementary abelian p -group, p and q are primes (not necessarily distinct), α is an automorphism of q -power order, and no nontrivial, proper subgroup of A is invariant under α .

Definition 6.7 ([4, p. 5]). Let G be a group. For any subgroups H and K of G , such that $K \triangleleft H$, the quotient H/K is said to be a *section* of G .

Theorem 6.8. Any non-CCA group of odd order has a section that is isomorphic to either:

1. a semi-wreathed product $A \wr_\alpha \mathbb{Z}_n$ (see Example 6.5), where A is a nontrivial, elementary abelian group (of odd order) and $n > 1$, or
2. the (unique) nonabelian group of order 21.

Proof. Assume $\text{Cay}(G; S)$ is a connected Cayley graph on a group G of odd order that does not have a section as described in either (1) or (2). We will show, by induction on the order, that if $\mathcal{A}$ is any subgroup of $\mathcal{A}^0$ that contains $\widehat{G}$, then $\widehat{G}$ is a normal subgroup of $\mathcal{A}$. (Then taking $\mathcal{A} = \mathcal{A}^0$ implies that $\text{Cay}(G; S)$ is CCA (see Remark 6.2).)

It is important to note that this conclusion implies $\widehat{G}$ is a *characteristic* subgroup of $\mathcal{A}$ (because Lemma 6.3 implies that $\widehat{G}$ is the unique largest normal subgroup of odd order). For convenience, we write $\widehat{G} \triangleleft \mathcal{A}$ when $\widehat{G}$ is characteristic.

Let $\mathcal{N}$ be a minimal normal subgroup of $\mathcal{A}$. Then $\mathcal{N}$ is either elementary abelian or the direct product of (isomorphic) nonabelian simple groups [4, Thm. 2.1.5, p. 17], and we consider the two possibilities as separate cases.

Case 1. Assume $\mathcal{N}$ is elementary abelian. Since the Sylow 2-subgroup $\mathcal{A}_e$, being the stabilizer of a vertex, does not contain any normal subgroups of $\mathcal{A}$, we know that $\mathcal{N}$ is not contained in a Sylow 2-subgroup. Hence, $\mathcal{N}$ is not a 2-group, so it must be a p -group for some odd prime p . Therefore, since $\widehat{G}$ is the largest normal subgroup of odd order, we have $\mathcal{N} \subseteq \widehat{G}$, so

$$\mathcal{N} = \widehat{N}, \text{ for some (elementary abelian) normal subgroup } N \text{ of } G.$$

Let $\mathcal{N}^+$ be the kernel of the action of $\mathcal{A}$ on G/N , so $\mathcal{A}/\mathcal{N}^+$ is a group of colour-preserving automorphisms of $\text{Cay}(G/N; \overline{S})$, where $\overline{S}$ is the image of S in G/N . Therefore, by induction on $|\mathcal{A}|$, we know that $\widehat{G}\mathcal{N}^+/\mathcal{N}^+$ is normal in $\mathcal{A}/\mathcal{N}^+$, so $\widehat{G}\mathcal{N}^+$ is normal in $\mathcal{A}$. Then we may assume $\widehat{G}\mathcal{N}^+ = \mathcal{A}$, for otherwise, by induction on $|\mathcal{A}|$, we would know $\widehat{G} \triangleleft \widehat{G}\mathcal{N}^+$, so $\widehat{G} \triangleleft \mathcal{A}$, as desired. Since $|G|$ is odd, this implies that $\mathcal{N}^+$ contains a Sylow 2-subgroup of $\mathcal{A}$. In fact, since $\mathcal{N}^+$ is normal and all Sylow 2-subgroups are conjugate, this implies that $\mathcal{N}^+$ contains every Sylow 2-subgroup. In particular, it contains $\mathcal{A}_e$. Therefore $\mathcal{N}^+ = \mathcal{N}\mathcal{A}_e$, so

$$\mathcal{N}\mathcal{A}_e \triangleleft \mathcal{A}.$$

This means that $\mathcal{A}_e$ acts trivially on G/N , so, for every $s \in S \setminus N$, $\mathcal{A}_e$ preserves the orientation of every s -edge. (This uses the fact that, since $|s|$ is odd, $s \not\equiv s^{-1} \pmod{N}$ if $s \notin N$.) This implies:

$$\text{for } \varphi \in \mathcal{A}_e, g \in G, \text{ and } x \in \langle S \setminus N \rangle, \text{ we have } \varphi(gx) = \varphi(g)x. \quad (6.9)$$

Let $(S \cap N)^{\langle S \setminus N \rangle} = \{gsg^{-1} \mid s \in S \cap N, g \in \langle S \setminus N \rangle\}$. Now, suppose $t \in (S \cap N)^{\langle S \setminus N \rangle}$ and $h \in N$. There exists $s \in S \cap N$ and $x \in \langle S \setminus N \rangle$, such that $xsx^{-1} = t$. From (6.9) and the fact that φ is colour-preserving, we see that

$$\varphi(ht) = \varphi(hxsx^{-1}) = \varphi(h)x s^{\pm 1} x^{-1} = \varphi(h)t^{\pm 1}.$$

Hence, $\varphi|_N$ is a colour-preserving automorphism of

$$\text{Cay}\left(N; (S \cap N)^{\langle S \setminus N \rangle} \cup (\langle S \setminus N \rangle \cap N)\right).$$

Since S generates G , it is easy to see that this Cayley graph is connected.

Note that $C_{\mathcal{A}_e}(\mathcal{N})$ is normalized by both $\mathcal{N}$ and $\mathcal{A}_e$, so it is a normal subgroup of $\mathcal{N}\mathcal{A}_e$. Therefore, it must be trivial (since the largest normal 2-subgroup of $\mathcal{N}\mathcal{A}_e$ is characteristic,

and is therefore normal in $\mathcal{A}$, but the stabilizer $\mathcal{A}_e$ does not contain any nontrivial normal subgroups of $\mathcal{A}$). So

$$\mathcal{A}_e \text{ acts faithfully by conjugation on } \mathcal{N}. \quad (6.10)$$

Also, we know that $\varphi|_N$ is an automorphism of N (by Remark 6.2, since φ normalizes $\mathcal{N} = \widehat{N}$). Since, being a colour-preserving automorphism, φ either centralizes or inverts every element of the generating set of N , this implies that $\varphi^2|_N$ is trivial. Since this is true for every $\varphi \in \mathcal{A}_e$, we conclude that $\mathcal{A}_e$ acts on N via an elementary abelian 2-group. From (6.10), we conclude that $\mathcal{A}_e$ is elementary abelian.

We can think of $\mathcal{N}$ as a vector space over $\mathbb{Z}_p$, and, for each homomorphism $\gamma: \mathcal{A} \rightarrow \{\pm 1\}$, let

$$\mathcal{N}_\gamma = \{n \in \mathcal{N} \mid ana^{-1} = \gamma(a)n \text{ for all } a \in \mathcal{A}_e\}.$$

(This is called the “weight space” associated to γ .) Since every linear transformation satisfying $T^2 = I$ is diagonalizable, and $\mathcal{A}_e$ is commutative, the elements of $\mathcal{A}_e$ can be simultaneously diagonalized. This means that if we let $\Gamma = \{\gamma \mid \mathcal{N}_\gamma \neq \{0\}\}$, then, since eigenspaces for different eigenvalues are always linearly independent, we have $\mathcal{N} = \bigoplus_{\gamma \in \Gamma} \mathcal{N}_\gamma$. This direct-sum decomposition is canonically defined from the action of $\mathcal{A}_e$ on $\mathcal{N}$. Since $\widehat{G}$ acts on $\mathcal{N}\mathcal{A}_e$ (by conjugation), we conclude that the action of $\widehat{G}$ on $\mathcal{N}$ by conjugation must permute the weight spaces. More precisely, there is an action of G on Γ , such that $\widehat{g}\mathcal{N}_\gamma\widehat{g}^{-1} = \mathcal{N}_{g\gamma}$ for all $g \in G$. Since N is abelian, this factors through to a well-defined action of G/N on Γ .

If the G -action on Γ is trivial, then every weight space is G -invariant, which implies that the action of $\widehat{G}$ on $\mathcal{N}$ commutes with the action of $\mathcal{A}_e$. Since $\mathcal{A}_e$ acts faithfully, we conclude that $\widehat{G}$ centralizes $\mathcal{A}_e\mathcal{N}/\mathcal{N}$; that is, $[\widehat{G}, \mathcal{A}_e] \subseteq \mathcal{N} \subseteq \widehat{G}$. So $\mathcal{A}_e$ normalizes $\widehat{G}$, as desired.

We may now assume that the G -action is nontrivial, so there is some $g \in G$ with an orbit of some length $n > 1$ on Γ . Let γ_0 be an element of this orbit, so $\widehat{g}^n$ normalizes $\mathcal{N}_{\gamma_0}$. Since $S \setminus N$ generates G/N , we may assume $g \in S \setminus N$, so (6.9) tells us that $\langle \widehat{g} \rangle \cap \mathcal{N}$ is centralized by $\mathcal{A}_e$. However, the minimality of $\mathcal{N}$ implies that $C_{\mathcal{N}}(\mathcal{A}_e) = \mathcal{N} \cap Z(\mathcal{N}\mathcal{A}_e)$ is trivial. Therefore, $\langle \mathcal{N}, \widehat{g} \rangle = \mathcal{N} \rtimes \langle \widehat{g} \rangle$ is a semidirect product. So

$$\langle \mathcal{N}_{\gamma_0}, \widehat{g} \rangle = \left(\bigoplus_{\gamma \in \langle g \rangle \gamma_0} \mathcal{N}_\gamma \right) \rtimes \langle \widehat{g} \rangle.$$

Then modding out $C_{\langle \widehat{g} \rangle}(\mathcal{N}_{\gamma_0})$ yields a section of $\widehat{G}$ that is isomorphic to $\mathcal{N}_{\gamma_0} \wr_{\alpha} \mathbb{Z}_n$, where α is the automorphism of $\mathcal{N}_{\gamma_0}$ induced by the conjugation action of $\widehat{g}^n$. So G has a semi-wreathed section, as described in (1). This completes the proof of this case.

Case 2. Assume $\mathcal{N} = \mathcal{L}_1 \times \cdots \times \mathcal{L}_r$, where each $\mathcal{L}_i$ is a nonabelian simple group, and $\mathcal{L}_i \cong \mathcal{L}_1$ for all i . We know that $\mathcal{A} = \widetilde{G}\mathcal{A}_e$, $\mathcal{A}_e$ is a 2-group, and $|G|$ is odd, so $\widehat{G}$ is a 2-complement in $\mathcal{A}$. (By definition, this means that $|\widehat{G}|$ is odd and $|\mathcal{A} : \widehat{G}|$ is a power of 2 [7, p. 88].) So $\mathcal{L}_1$ is a nonabelian simple group that has a 2-complement (namely, $\widehat{G} \cap \mathcal{L}_1$). By using the Classification of Finite Simple Groups, it can be shown that this implies $\mathcal{L}_1 \cong \text{PSL}(2, p)$, for some Mersenne prime $p \geq 7$ (see [9, Thm. 1.3]).

Note that $\mathcal{A}_e \cap \mathcal{L}_i$ is a Sylow 2-subgroup of $\mathcal{L}_i \cong \text{PSL}(2, p)$. Therefore, it is dihedral [4, Lem. 15.1.1(iii)] and has order $p + 1$ (because p is a Mersenne prime). Let

- $\mathcal{C}_i$ be the unique cyclic subgroup of order $(p+1)/2$ in $\mathcal{A}_e \cap \mathcal{L}_i$,
- $\mathcal{C}_i^2$ be the unique subgroup of index 2 in $\mathcal{C}_i$, and
- $\mathcal{C}^2 = \mathcal{C}_1^2 \times \cdots \times \mathcal{C}_r^2 \subset \mathcal{A}_e \cap (\mathcal{L}_1 \times \cdots \times \mathcal{L}_r)$.

Since every element of $\mathcal{C}_i$ is a colour-preserving automorphism, it either fixes or inverts each element of S , so we know that $\mathcal{C}_i^2$ fixes every element of S . Since stabilizers are conjugate, this implies $\widehat{s}^{-1}\mathcal{C}_i^2\widehat{s} \subseteq \mathcal{A}_e$, for every $s \in S$. We must have $p > 7$, for otherwise $\widehat{G} \cap \mathcal{L}_i$, being the 2-complement of $\text{PSL}(2, 7)$, would be the nonabelian group of order 21, as in (2). This implies that $\mathcal{C}_i^2$ is the unique cyclic subgroup of order $(p+1)/4$ in the dihedral group $\mathcal{A}_e \cap \mathcal{L}_i$, so we must have $\widehat{s}^{-1}\mathcal{C}_i^2\widehat{s} = \mathcal{C}_i^2$, which means that $\widehat{s}$ normalizes $\mathcal{C}_i^2$. Since this holds for every s in the generating set S , we conclude that $\widehat{G}$ normalizes $\mathcal{C}^2$.

Note that $\mathcal{A}_e$ normalizes $\mathcal{A}_e \cap \mathcal{N}$, and that $\mathcal{C}^2 \triangleleft \mathcal{A}_e \cap \mathcal{N}$ (since, as was mentioned above, $\mathcal{C}_i$ is the unique cyclic subgroup of its order in $\mathcal{A}_e \cap \mathcal{L}_i$). Therefore, $\mathcal{C}^2 \triangleleft \mathcal{A}_e$. We conclude that $\mathcal{C}^2$ is normal in $\widehat{G}\mathcal{A}_e = \mathcal{A}$. So $\mathcal{C}_1^2 = \mathcal{C}^2 \cap \mathcal{L}_1$ is normal in $\mathcal{L}_1$, contradicting the fact that $\mathcal{L}_1$ is simple. $\square$

Lemma 6.11. A group G is strongly CCA (or CCA) if and only if, for every generating set S of G such that every element of S has prime-power order, the Cayley graph $\text{Cay}(G; S)$ is strongly CCA (or CCA).

Proof. Suppose φ is a colour-permuting automorphism of some connected Cayley graph $\text{Cay}(G; S)$. There is a permutation π of S , such that $\varphi(gs) = \varphi(g)\pi(s)^{\pm 1}$, for all $g \in G$ and $s \in S$. (Furthermore, if φ is colour-preserving, then π can be taken to be the identity permutation.) By induction on k , this implies $\varphi(gs^k) = \varphi(g)\pi(s)^{\pm k}$, for all $k \in \mathbb{Z}$. Hence, if we let $S^* = \{s^k \mid s \in S, k \in \mathbb{Z}\}$, then φ is a colour-permuting automorphism of $\text{Cay}(G; S^*)$. Now, let

$$S_0 = \{t \in S^* \mid |t| \text{ is a prime-power}\}.$$

Then φ is a colour-permuting automorphism of $\text{Cay}(G; S_0)$, and S_0 generates G , since every element s of the generating set S can be written as a product of elements of $\langle s \rangle$ that have prime-power order [4, Thm. 1.3.1(iii), p. 9], and therefore belong to S_0 . (Furthermore, φ is colour-preserving if the permutation π is the identity permutation.) $\square$

Lemma 6.12. Suppose

- C is a cyclic, normal subgroup of a group H ,
- $|C|$ is relatively prime to $|H : C|$,
- no element of $H \setminus C$ centralizes C , and
- α is an automorphism of H .

Then $\alpha(h) \in hC$, for every $h \in H$.

Proof. Since no other subgroup of H has the same order as C , we know that $\alpha|_C$ is an automorphism of C , so there exists $r \in \mathbb{Z}$, such that $\alpha(c) = c^r$, for every $c \in C$. Then, for every $h \in H$ and $c \in C$, we have

$$\alpha(h) c^r \alpha(h)^{-1} = \alpha(h) \alpha(c) \alpha(h)^{-1} = \alpha(hch^{-1}) = (hch^{-1})^r = hc^r h^{-1},$$

so $h^{-1}\alpha(h)$ centralizes C . By assumption, this implies $h^{-1}\alpha(h) \in C$, as desired. $\square$

Corollary 6.13. The following are equivalent:

1. There is a group of order n that is not CCA.
2. There is a group of order n that is not strongly CCA.
3. $n \geq 8$, and n is divisible by either 4, 21, or a number of the form $p^q \cdot q$, where p and q are primes (not necessarily distinct) and p is odd.

Proof. (1 $\Rightarrow$ 2) Obvious.

(3 $\Rightarrow$ 1) If n is divisible by 4, then there is a generalized dicyclic group of order n , which is not CCA (see Corollary 2.8(2)). The nonabelian group of order 21 and the wreath product $\mathbb{Z}_p \wr \mathbb{Z}_q$ (which is of order $p^q \cdot q$) are not CCA (see Examples 2.3 and 2.4). Taking an appropriate direct product yields a non-CCA group whose order is any multiple of these (see Proposition 3.1).

(2 $\Rightarrow$ 3) Assume there is a group G of order n that is not strongly CCA, but n is not divisible by 4, 21, or a number of the form $p^q \cdot q$. From Theorem 6.8, we see that n is even. (Otherwise, $n = |G|$ is divisible by the order of a semi-wreathed product $|A \wr \mathbb{Z}_k|$. If we let p and q be prime divisors of $|A|$ and k , respectively, then $|A \wr \mathbb{Z}_k| = |A|^k \cdot k$ is a multiple of $p^q \cdot q$.) Furthermore, n must be square-free, for otherwise it is a multiple of either 4 or $p^2 \cdot 2$, for some prime p . Therefore, G is a semidirect product $\mathbb{Z}_k \rtimes \mathbb{Z}_\ell$ [6, Cor. 9.4.1].

We may assume the centre of G is trivial, for otherwise we can write G as a nontrivial direct product, so Proposition 3.2 (and induction on n) implies that G is CCA. Therefore, k is odd (so ℓ is even), so we may write $G = \mathbb{Z}_k \rtimes (\mathbb{Z}_m \times \mathbb{Z}_2)$, and $\mathbb{Z}_m \times \mathbb{Z}_2$ acts faithfully on $\mathbb{Z}_k$. Let $H = \mathbb{Z}_k \rtimes \mathbb{Z}_m$, so $|H| = km$ is odd, and H is the (unique) subgroup of index 2 in G .

Let φ be a colour-permuting automorphism of a connected Cayley graph $\text{Cay}(G; S)$. (We wish to show that φ is affine.) There is no harm in assuming that every element of S has prime order (see Lemma 6.11).

Case 1. Assume φ is colour-preserving. Fix some $t \in S$ with $|t| = 2$. We claim we may assume that t is the only element of order 2 in S , and that $H = \langle S \setminus \{t\} \rangle$. To see this, let

- T be the set of all elements of order 2 in S , and
- $S' = \{t\} \cup \{uv \mid u, v \in T, u \neq v\} \cup (S \setminus T)$.

It is easy to see that φ is a colour-preserving automorphism of the connected Cayley graph $\text{Cay}(G; S')$, and that $G = \langle S \setminus \{t\} \rangle \langle t \rangle$. This establishes the claims.

From Theorem 6.8 (and the fact that $|H|$ is odd), we know that $\varphi|_H$ is affine. By composing with a left translation, we may assume that φ fixes e . Then $\varphi|_H$ is a group automorphism. By composing with an automorphism of $\mathbb{Z}_k \rtimes (\mathbb{Z}_m \times \mathbb{Z}_2)$ of the form $(x, y, z) \mapsto (x^r, y, z)$, we may assume $\varphi|_{\mathbb{Z}_k}$ is the identity map. Also, since $\varphi(s) \in \{s^{\pm 1}\}$ for every $s \in S$, and $|H/\mathbb{Z}_k| = m$ is odd, Lemma 6.12 implies that φ also fixes every element of $(S \cap H) \setminus \mathbb{Z}_k$. Hence, $\varphi|_H$ is an automorphism that fixes every element of a generating set, so $\varphi(h) = h$ for every $h \in H$. Since $\varphi(ht) = \varphi(h)t = ht$, for all $h \in H$ (because φ is colour-preserving and $t = t^{-1}$), we conclude that φ fixes every element of G , and is therefore affine, as desired.

Case 2. The general case. From Case 1, we see that G is CCA, so $\widehat{G} \triangleleft \mathcal{A}^0$. Hence, $\widehat{H} \triangleleft \mathcal{A}^0$ (since it is the unique largest normal subgroup of odd order), so φ normalizes $\widehat{H}$. This implies that the restriction of φ to H is an automorphism of H .

For each $s \in S$, let $\tilde{s} = \varphi(s) \in S$. To prove that φ is affine, it suffices to show $\varphi(xs) = \varphi(x)\tilde{s}$ for all $x \in G$ and $s \in S$ (see Remark 1.6(4)). If this is not the case, then, since φ is colour-permuting, there must be some x , such that $\varphi(xs) = \varphi(x)\tilde{s}^{-1}$ (and $\tilde{s}^{-1} \neq \tilde{s}$, which means $|s| \neq 2$). This will lead to a contradiction.

Since $|s| \neq 2$, and we have assumed that every element of S has prime order (by Lemma 6.11), we see that $s \in H$. Then, since $\varphi|_H$ is an automorphism, but

$$\varphi(xs) = \varphi(x)\tilde{s}^{-1} = \varphi(x)\varphi(s)^{-1} \neq \varphi(x)\varphi(s),$$

we must have $x \notin H$. Since H has only two cosets, and there must be some element of S that is not in H , this implies that we may assume $x \in S$, after multiplying on the left by an appropriate element of H (and using the fact that φ normalizes $\tilde{H}$). Note that, since $x \notin H$, and every element of S has prime order, this implies $|x| = 2$. So the order of $\tilde{x}$ is also 2, which implies $\tilde{x} \notin H$ (since $|H|$ is odd).

Since φ is colour-permuting, we have

$$\varphi(xs) = \varphi({}^x s x) = \varphi({}^x s)\tilde{x}.$$

Also, by the choice of x and s , we have

$$\varphi(xs) = \varphi(x)\tilde{s}^{-1} = \tilde{x}\tilde{s}^{-1}.$$

Therefore

$$\varphi({}^x s) = \tilde{x}\tilde{s}^{-1}.$$

Since $\mathbb{Z}_m$ acts faithfully on $\mathbb{Z}_k$, we have $\alpha(h) \equiv h \pmod{\mathbb{Z}_k}$, for every automorphism α of H (see Lemma 6.12). Since φ and conjugation by x are automorphisms of H , this implies $s \equiv s^{-1} \pmod{\mathbb{Z}_k}$. Since $|s|$ is odd, we conclude that $s \in \mathbb{Z}_k$.

Then, since the automorphism group of a cyclic group is abelian, we have

$$\varphi({}^x s) = {}^x \varphi(s) = {}^x \tilde{s},$$

so $x^{-1}\tilde{x}$ must invert $\tilde{s}$. But this is impossible, because, as was mentioned above, x and $\tilde{x}$, being of order 2, cannot be in H , so they are both in the other coset of H , so $x^{-1}\tilde{x} \in H$ has odd order. This contradiction completes the proof that φ is affine. $\square$

Remark 6.14. It is not necessary to assume p is odd in the statement of Corollary 6.13(3), because $2^q \cdot q$ is divisible by 4, which is already in the list of divisors.

7 Groups of small order

In this section, we briefly explain which groups of order less than 32 are CCA (or strongly CCA). First, note that almost all of the abelian ones are strongly CCA:

Proposition 7.1 (cf. Proposition 4.1). An abelian group of order less than 32 is not strongly CCA if and only if it is either

- $\mathbb{Z}_2 \times \mathbb{Z}_4$ (of order 8),
- $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_4$ (of order 16), or
- $\mathbb{Z}_2 \times \mathbb{Z}_3 \times \mathbb{Z}_4$ (of order 24).

None of these are CCA.

Also note that almost all of the groups whose order is not divisible by 4 are CCA:

Proposition 7.2. The only groups that are not strongly CCA, and whose order is less than 32 and not divisible by 4 are:

- the wreath product $\mathbb{Z}_3 \wr \mathbb{Z}_2$, which is isomorphic to $D_6 \times \mathbb{Z}_3$ and has order 18, and
- the nonabelian group of order 21.

Neither of these is CCA.

Proof. For the groups of odd order, the conclusion is immediate from Theorem 6.8 and Example 2.3 (see Proposition 7.4 for a stronger result). Proposition 3.2 deals with the groups $D_6 \times \mathbb{Z}_5$ and $D_{10} \times \mathbb{Z}_3$ of order 30. For all of the other groups of even order, it suffices to note that if m is odd, then every generalized dihedral group of order $2m$ is strongly CCA (see Proposition 5.6). $\square$

So it is surprising that very few of the remaining groups are strongly CCA:

Proposition 7.3. The only nonabelian groups that are strongly CCA and whose order is less than 32 and divisible by 4 are:

- the dihedral groups of order 8, 16, and 24,
- the alternating group A_4 , which is of order 12,
- another group of order 16, namely, the semidirect product

$$\mathbb{Z}_8 \rtimes \mathbb{Z}_2 = \langle x, a \mid x^8 = a^2 = e, a^{-1}xa = x^5 \rangle,$$

and

- three additional groups of order 24, namely, $D_8 \times \mathbb{Z}_3$, $A_4 \times \mathbb{Z}_2$, and the semidirect product $\mathbb{Z}_3 \rtimes \mathbb{Z}_8$ in which $\mathbb{Z}_8$ inverts $\mathbb{Z}_3$.

Furthermore, the only groups of order less than 32 that are CCA, but not strongly CCA, are:

- the dihedral groups D_{12} , D_{20} , and D_{28} , and
- the group $D_{12} \times \mathbb{Z}_2$, which is a generalized dihedral group of order 24.

Sketch of proof. The result can be verified by an exhaustive computer search, but we summarize a case-by-case analysis that can be carried out by hand, using the classification of groups of order less than 32. Each group of such small order can be specified by its “GAP Id,” which is an ordered pair $[n, k]$, where n is the order of the group, and k is the id number that has been assigned to that particular group (see [5], for example).

Assume G is nonabelian, $|G| < 32$, and $|G|$ is divisible by 4. We may assume that G is neither generalized dicyclic, semidihedral, nor generalized dihedral, for otherwise Corollary 2.8(2,3) and Propositions 5.3 and 5.6 determine whether G is CCA or strongly CCA. By inspection of the list of groups of each order, we see that this leaves only thirteen possibilities for G , and we consider each of these GAP Ids separately. In most cases, Proposition 2.5 implies that G is not CCA.

See Appendix A for some additional details.

See Ap-
pendix B
for some
additional
details.

See Ap-
pendix B
for some
additional
details.

See Ap-
pendix B
for some
additional
details.

See Ap-
pendix B
for some
additional
details.

- [12, 3] = A_4 . This group is strongly CCA, but we omit the proof.
- [16, 3] = $\langle a, b, c \mid a^4 = b^2 = c^2 = e, ab = ba, bc = cb, cac = ab \rangle$. Proposition 2.5 applies with $S = \{a^{\pm 1}, c\}$, $T = \{a^{\pm 1}\}$, and $\tau = a^2 \in Z(G)$.
- [16, 6] = $\langle a, x \mid a^8 = x^2 = e, xax = a^5 \rangle = \langle a \rangle \rtimes \langle x \rangle = \mathbb{Z}_8 \rtimes \mathbb{Z}_2$. This group is strongly CCA, but we omit the proof.
- [16, 13] = $\langle a, x, y \mid a^4 = x^2 = e, a^2 = y^2, xax = a^{-1}, ay = ya, xy = yx \rangle$. Proposition 2.5 applies with $S = \{a^{\pm 1}, x, y^{\pm 1}\}$, $T = \{a^{\pm 1}, y^{\pm 1}\}$, and $\tau = a^2 \in Z(G)$.
- [20, 3] = $\langle a, b \mid a^5 = b^4 = e, bab^{-1} = a^2 \rangle$. Proposition 2.5 applies with $S = \{a^{\pm 1}, b^{\pm 1}\}$, $T = \{b^{\pm 1}\}$, and $\tau = b^2$ (which inverts a).
- [24, 1] = $\mathbb{Z}_3 \rtimes \mathbb{Z}_8$, where $\mathbb{Z}_8$ inverts $\mathbb{Z}_3$. This group is strongly CCA, but we omit the proof.
- [24, 3] = $\text{SL}(2, 3) \cong Q_8 \rtimes \mathbb{Z}_3 = \langle i, j \rangle \rtimes \langle a \rangle$, where $aia^{-1} = j$ and $a^{-1}ia = ij$. Proposition 2.5 applies with $S = \{i^{\pm 1}, a^{\pm 1}\}$, $T = \{i^{\pm 1}\}$, and $\tau = i^2 \in Z(G)$.
- [24, 5] = $S_3 \times \mathbb{Z}_4$. Proposition 2.5 applies with $T = \{(1, 2)\} \times \{\pm 1\}$, $S = \{((2, 3), 0)\} \cup T$, and $\tau = (e, 2) \in Z(G)$.
- [24, 8] = $\mathbb{Z}_3 \rtimes D_8 = \langle a, b, c \mid a^3 = b^4 = c^2 = e, bab^{-1} = a^{-1}, ac = ca, cbc^{-1} = b^{-1} \rangle$. Proposition 2.5 applies with $S = \{(ab)^{\pm 1}, b^{\pm 1}, c\}$, $T = \{(ab)^{\pm 1}, b^{\pm 1}\}$, and $\tau = b^2 \in Z(G)$.
- [24, 10] = $D_8 \times \mathbb{Z}_3$. Since D_8 is strongly CCA (see Proposition 5.6), the same is true for this group (see Proposition 3.2).
- [24, 11] = $Q_8 \times \mathbb{Z}_3$. This is not CCA, since Q_8 is not CCA (see Example 2.2 and Proposition 3.1).
- [24, 12] = S_4 . Let $a = (1, 2, 3, 4)$ and $b = (1, 2, 4, 3)$, so Proposition 2.5 applies, with $S = \{a^{\pm 1}, b^{\pm 1}\}$, $T = \{a^{\pm 1}\}$, and $\tau = a^2 = (1, 3)(2, 4)$, which inverts b .
- [24, 13] = $A_4 \times \mathbb{Z}_2$. This group is strongly CCA, but we omit the proof. $\square$

The above results assume $|G| < 32$, but it is not difficult to treat considerably larger groups if we assume the order is odd:

Proposition 7.4. Let $G_{21} = \mathbb{Z}_7 \rtimes \mathbb{Z}_3$ be the (unique) nonabelian group of order 21. Then the only groups of odd order less than 100 that are not strongly CCA are G_{21} , $G_{21} \times \mathbb{Z}_3$, and $\mathbb{Z}_3 \wr \mathbb{Z}_3$.

Proof. Suppose G is a group of odd order, such that G is not strongly CCA and $|G| < 100$. From Corollary 6.13, we see that $|G|$ is divisible by either 21 or $3^3 \cdot 3 = 81$. Since $|G| < 100$, this implies that $|G|$ is either 21, $21 \times 3 = 63$, or $3^3 \cdot 3 = 81$. Also, G must be nonabelian (see Corollary 4.2).

- The nonabelian group G_{21} of order 21 is not CCA (see Example 2.3).
- There are two nonabelian groups of order 63. One of them, the direct product $G_{21} \times \mathbb{Z}_3$, is not CCA (see Proposition 3.1). The other is

$$\mathbb{Z}_7 \rtimes \mathbb{Z}_9 = \langle x, a \mid x^7 = a^9 = e, a^{-1}xa = x^2 \rangle.$$

This group is strongly CCA, but we omit the proof.

- 6.8 implies that $\mathbb{Z}_3 \wr \mathbb{Z}_3$ is the only non-CCA group of order 81 (see also Example 2.4). $\square$

See Ap-
pendix B
for some
additional
details.

References

- [1] M. Albert, J. Bratz, P. Cahn, T. Fargus, N. Haber, E. McMahon, J. Smith and S. Tekansik, Color-permuting automorphisms of Cayley graphs. *Congr. Numer.* **190** (2008), 161–171. MR 2489799
- [2] E. Dobson, Some non-normal Cayley digraphs of the generalized quaternion group of certain orders, *Electron. J. Combin.* **10** (2003), #R31, 7 pages. MR 2014518,
- [3] M. L. Fiol, M. A. Fiol and J. L. A. Yebra, When the arc-colored line digraph of a Cayley colored digraph is again a Cayley colored digraph, *Ars Combin.* **34** (1992), 65–73. MR 1206550
- [4] D. Gorenstein, *Finite Groups*, second ed., Chelsea, New York, 1980. ISBN 0-8284-0301-5, MR 0569209
- [5] Groupprops, The Group Properties Wiki: Groups of a particular order, http://groupprops.subwiki.org/wiki/Category:Groups_of_a_particular_order
- [6] M. Hall, *The Theory of Groups*, Chelsea, New York, 1976. ISBN 978-0-8218-1967-8, MR 0414669
- [7] I. M. Isaacs, *Finite Group Theory*, American Mathematical Society, Providence, RI, 2008. ISBN 978-0-8218-4344-4, MR 2426855
- [8] C. H. Li, On isomorphisms of finite Cayley graphs—a survey, *Discrete Math.* **256** (2002) 301–334. MR 1927074
- [9] C. Martinez-Perez and W. Willems, The trivial intersection problem for characters of principal indecomposable modules, *Adv. Math.* **222** (2009), 1197–1219. MR 2554934
- [10] J. Morris, Automorphisms of circulants that respect partitions (preprint). <http://arxiv.org/abs/1205.0503>
- [11] S. K. Sehgal, On the normalizer of a group in the Cayley representation, *Internat. J. Math. Math. Sci.* **12** (1989), 459–462. MR 1007196
- [12] A. T. White, *Graphs, Groups and Surfaces*, Elsevier, New York, 1973. MR 0340026
- [13] M.–Y. Xu, Automorphism groups and isomorphisms of Cayley digraphs, *Discrete Math.* **182** (1998), 309–319. MR 1603719

APPENDIX: Notes to aid the referee

A Dicyclic (and other) groups omitted from the proof of Proposition 7.3

In order to keep the proof of Proposition 7.3 short, it has no discussion of groups that are generalized dicyclic, semidihedral, or generalized dihedral. Here are additional details to show that no cases were missed.

order 4: http://groupprops.subwiki.org/wiki/Groups_of_order_4
There are no nonabelian groups of order 4.

order 8: http://groupprops.subwiki.org/wiki/Groups_of_order_8
[8, 1] = $\mathbb{Z}_8$ is abelian
[8, 2] = $\mathbb{Z}_4 \times \mathbb{Z}_2$ is abelian
[8, 3] = D_8 is dihedral (and is **strongly CCA**, since $\mathbb{Z}_4$ does not have $\mathbb{Z}_2$ as a direct factor)
[8, 4] = Q_8 is dicyclic (so it is not CCA)
[8, 5] = $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$ is abelian

order 12: http://groupprops.subwiki.org/wiki/Groups_of_order_12
[12, 1] = $Q_{12} = \text{Dic}(3, \mathbb{Z}_6)$ is dicyclic (so it is not CCA)
[12, 2] = $\mathbb{Z}_{12}$ is abelian
[12, 3] = A_4 is discussed in the proof of Proposition 7.3. It is **strongly CCA**.
[12, 4] = D_{12} is dihedral (and is **CCA but not strongly CCA**, since $\mathbb{Z}_6$ has $\mathbb{Z}_2$ as a direct factor)
[12, 5] = $\mathbb{Z}_6 \times \mathbb{Z}_2$ is abelian

order 16: http://groupprops.subwiki.org/wiki/Groups_of_order_16
[16, 1] = $\mathbb{Z}_{16}$ is abelian
[16, 2] = $\mathbb{Z}_4 \times \mathbb{Z}_4$ is abelian
[16, 3] is discussed in the proof of Proposition 7.3. It is not CCA.
[16, 4] = $\mathbb{Z}_4 \rtimes \mathbb{Z}_4 = \text{Dic}((0, 1), \mathbb{Z}_4 \times \mathbb{Z}_2)$ is generalized dicyclic (so it is not CCA).
[16, 5] = $\mathbb{Z}_8 \times \mathbb{Z}_2$ is abelian
[16, 6] is discussed in the proof of Proposition 7.3. It is **strongly CCA**.
[16, 7] = D_{16} is dihedral (and is **strongly CCA**, since $\mathbb{Z}_8$ does not have $\mathbb{Z}_2$ as a direct factor)
[16, 8] is semidihedral (so it is not CCA).
[16, 9] = $Q_{16} = \text{Dic}(4, \mathbb{Z}_8)$ is dicyclic (so it is not CCA)
[16, 10] = $\mathbb{Z}_4 \times \mathbb{Z}_2 \times \mathbb{Z}_2$ is abelian
[16, 11] = $D_8 \times \mathbb{Z}_2$ is generalized dihedral over $\mathbb{Z}_4 \times \mathbb{Z}_2$ (and it is not CCA, since $\mathbb{Z}_4 \times \mathbb{Z}_2$ is not CCA).
[16, 12] = $Q_8 \times \mathbb{Z}_2 = \text{Dic}((2, 0), \mathbb{Z}_4 \times \mathbb{Z}_2)$ is generalized dicyclic (so it is not CCA).
[16, 13] is discussed in the proof of Proposition 7.3. It is not CCA.
[16, 14] = $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$ is abelian

order 20: http://groupprops.subwiki.org/wiki/Groups_of_order_20

[20, 1] = $Q_{20} = \text{Dic}(5, \mathbb{Z}_{10})$ is dicyclic (so it is not CCA)

[20, 2] = $\mathbb{Z}_{20}$ is abelian

[20, 3] is discussed in the proof of Proposition 7.3. It is not CCA.

[20, 4] = D_{20} is dihedral (and is **CCA but not strongly CCA**, since $\mathbb{Z}_{10}$ has $\mathbb{Z}_2$ as a direct factor)

[20, 5] = $\mathbb{Z}_{10} \times \mathbb{Z}_2$ is abelian

order 24: http://groupprops.subwiki.org/wiki/Groups_of_order_24

[24, 1] is discussed in the proof of Proposition 7.3. It is **strongly CCA**.

[24, 2] = $\mathbb{Z}_{24}$ is abelian

[24, 3] is discussed in the proof of Proposition 7.3. It is not CCA.

[24, 4] = $Q_{24} = \text{Dic}(6, \mathbb{Z}_{12})$ is dicyclic (so it is not CCA).

[24, 5] is discussed in the proof of Proposition 7.3. It is not CCA.

[24, 6] = D_{24} is dihedral (and is **strongly CCA**, since $\mathbb{Z}_{12}$ does not have $\mathbb{Z}_2$ as a direct factor)

[24, 7] = $Q_{12} \times \mathbb{Z}_2 = \text{Dic}((3, 0), \mathbb{Z}_6 \times \mathbb{Z}_2)$ is generalized dicyclic (so it is not CCA).

[24, 8] is discussed in the proof of Proposition 7.3. It is not CCA.

[24, 9] = $\mathbb{Z}_6 \times \mathbb{Z}_4$ is abelian

[24, 10] is discussed in the proof of Proposition 7.3. It is **strongly CCA**.

[24, 11] is discussed in the proof of Proposition 7.3. It is not CCA.

[24, 12] is discussed in the proof of Proposition 7.3. It is not CCA.

[24, 13] is discussed in the proof of Proposition 7.3. It is **strongly CCA**.

[24, 14] = $D_{12} \times \mathbb{Z}_2$ is generalized dihedral over $\mathbb{Z}_6 \times \mathbb{Z}_2$ (and is **CCA but not strongly CCA**, since $\mathbb{Z}_6 \times \mathbb{Z}_2$ obviously has $\mathbb{Z}_2$ as a direct factor)

[24, 15] = $\mathbb{Z}_3 \times \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$ is abelian

order 28: There does not seem to be a web page for the groups of this order, but it is easy to classify the nonabelian ones, by noting that Sylow's Theorem implies the Sylow 7-subgroup is normal. This implies $G = \mathbb{Z}_7 \rtimes H$, where $|H| = 4$ (so H is either $\mathbb{Z}_4$ or $\mathbb{Z}_2 \times \mathbb{Z}_2$).

If $H = \mathbb{Z}_4$, then the only nonabelian semidirect product is the dicyclic group $Q_{28} = \text{Dic}(7, \mathbb{Z}_{14})$ (which is not CCA).

If $H = \mathbb{Z}_2 \times \mathbb{Z}_2$, then the only nonabelian semidirect product is the dihedral group D_{28} . (It is **CCA, but not strongly CCA**, since $\mathbb{Z}_{14}$ has $\mathbb{Z}_2$ as a direct factor.)

B Strongly CCA groups omitted from the proof of Proposition 7.3

This appendix proves that certain small groups (listed in Examples B.2 and B.3) are strongly CCA. These verifications were omitted from our proofs of Propositions 7.3 and 7.4. The following simple observation will play a key role.

Lemma B.1. Let

- φ be a colour-permuting automorphism of a Cayley graph $\text{Cay}(G; S)$, such that $\varphi(e) = e$,
- $\tilde{a} = \varphi(a)$ and $\tilde{b} = \varphi(b)$, for some $a, b \in S$,
- $\tau(v) \in \{\pm 1\}$, such that $\varphi(va) = \varphi(v) \tilde{a}^{\tau(v)}$, for all $v \in G$, and
- $k_1, k_2, \dots, k_{2r} \in \mathbb{Z} \setminus \{0\}$, such that $a^{k_1} b^{k_2} a^{k_3} \dots b^{k_{2r}} = e$ (and $r \geq 2$).

If $\epsilon_1 = \epsilon_3$ and $\epsilon_2 = \epsilon_4$, for all $\epsilon_1, \dots, \epsilon_{2r} \in \{\pm 1\}$, such that $\tilde{a}^{\epsilon_1 k_1} \tilde{b}^{\epsilon_2 k_2} \dots \tilde{b}^{\epsilon_{2r} k_{2r}} = e$, then $\varphi(va) = \varphi(v) \tilde{a}$ and $\varphi(vb) = \varphi(v) \tilde{b}$, for all $v \in \langle a^{k_3}, b^{k_2} \rangle$.

Proof. Since φ is colour-permuting, there exist $\sigma, \tau: G \rightarrow \{\pm 1\}$, such that

$$\varphi(va) = \varphi(v) \tilde{a}^{\sigma(v)} \text{ and } \varphi(vb) = \varphi(v) \tilde{b}^{\tau(v)} \text{ for all } v \in G.$$

We wish to show $\sigma(v) = \tau(v) = 1$ for all $v \in \langle a^{k_3}, b^{k_2} \rangle$. Since $\sigma(e) = \tau(e) = 1$, it suffices to show that $\sigma(vb^{k_2}) = \tau(va^{k_3}) = \tau(v)$ for all $v \in G$.

The two parts of the proof are very similar, so we show only that $\sigma(vb^{k_2}) = \sigma(v)$. The relation $a^{k_1} b^{k_2} a^{k_3} \dots b^{k_{2r}} = e$ represents a closed walk starting at v (or at any other desired vertex). Applying φ yields a closed walk starting at $\varphi(v)$. Since φ is colour-permuting, this closed walk corresponds to a relation of the form $\tilde{a}^{\epsilon_1 k_1} \tilde{b}^{\epsilon_2 k_2} \dots \tilde{b}^{\epsilon_{2r} k_{2r}} = e$, with $\epsilon_i \in \{\pm 1\}$. By assumption, we must have $\epsilon_1 = \epsilon_3$. Therefore

$$\sigma(a^{k_1} b^{k_2}) = \epsilon_3 = \epsilon_1 = \sigma(v).$$

This establishes the desired conclusion, since $\sigma(v) = \sigma(va^{k_1})$, and va^{k_1} is an arbitrary element of G . $\square$

Example B.2. The groups $[12, 3]$, $[16, 6]$, $[24, 1]$, and $[24, 13]$ from the proof of Proposition 7.3 are strongly CCA.

Proof. We consider each of the four groups individually; for convenience, let G be the group under consideration. Suppose φ is a colour-permuting automorphism of a connected Cayley graph $\text{Cay}(G; S)$, such that $\varphi(e) = e$, and let $\tilde{s} = \varphi(s)$, for each $s \in S$. We wish to show $\varphi \in \text{Aut } G$.

Assume $G = [12, 3]$. Let $a \in S$ with $|a| = 3$, and let N be the (unique) subgroup of order 4 in G .

Assume, for the moment, that there exists $b \in S \cap N$ (so $|b| = 2$). Then $(ab)^3 = e$. Suppose $i, j, k \in \{\pm 1\}$, with

$$e = \tilde{a}^i \tilde{b} \tilde{a}^j \tilde{b} \tilde{a}^k \tilde{b} \equiv \tilde{a}^{i+j+k} \pmod{N},$$

so $i+j+k \equiv 0 \pmod{3}$. Since $i, j, k \in \{\pm 1\}$, this implies $i = j = k$. We conclude from Lemma B.1 that $\varphi(vs) = \varphi(v) \tilde{s}$, for all $v \in \langle a, b \rangle = G$ and $s \in \{a^{\pm 1}, b\}$, so $\varphi \in \text{Aut } G$.

We may now assume $|s| = 3$ for all $s \in S$. Let $b \in S \setminus \langle a \rangle$. We may assume $a \equiv b \pmod{N}$, by replacing b with its inverse if necessary. Write $\tilde{b} = \tilde{a}^r x$, with $r \in \{\pm 1\}$ and $x \in N$. Note that $(a^{-1}b)^2 = e$. Suppose $i, j, k, \ell \in \{\pm 1\}$, with

$$e = \tilde{a}^{-i} \tilde{b}^j \tilde{a}^{-k} \tilde{b}^\ell = \tilde{a}^{-i+rj-k+r\ell} \cdot \begin{cases} (\tilde{a}^{k-r\ell} x \tilde{a}^{-k+r\ell})x & \text{if } j = \ell = 1, \\ (\tilde{a}^k x \tilde{a}^{-k})x & \text{if } j = -1 \text{ and } \ell = 1, \\ \tilde{a}^{-r\ell} (\tilde{a}^k x \tilde{a}^{-k})x \tilde{a}^{r\ell} & \text{if } j = 1 \text{ and } \ell = -1, \\ \tilde{a}^{-r\ell} (\tilde{a}^{k-rj} x \tilde{a}^{-k+rj})x \tilde{a}^{r\ell} & \text{if } j = \ell = -1. \end{cases}$$

Since the component in N must be trivial, and no nontrivial power of $\tilde{a}$ centralizes x , we see that we must have $j = \ell$ and $k = rj = r\ell$. Then, since the exponent of $\tilde{a}$ must be 0, this implies $i = k$. We conclude from Lemma B.1 that $\varphi(vs) = \varphi(v)\tilde{s}$, for all $v \in \langle a, b \rangle = G$ and $s \in \{a^{\pm 1}, b^{\pm 1}\}$, so $\varphi \in \text{Aut } G$.

Assume $G = [16, 6]$. Let $a \in S$ with $|a| = 8$. Let $b \in S \setminus \langle a \rangle$.

Assume, for the moment, that $|b| = 8$. Write $b^2 = a^{2r}$, for some odd r . Then we must have $\tilde{b}^2 = \tilde{a}^{2r}$. This implies that if $i, j \in \{\pm 1\}$, such that

$$e = \tilde{b}^{2i} \tilde{a}^{-2rj},$$

then $i = j$ (since $|\tilde{b}^2| = |b^2| > 2$). We conclude (much as in Lemma B.1) that $\varphi(vs) = \varphi(v)\tilde{s}$, for all $v \in \langle a, b \rangle = G$ and $s \in \{a^{\pm 1}, b^{\pm 1}\}$, so $\varphi \in \text{Aut } G$.

We may now assume $|b| \in \{2, 4\}$, so $b^2 \in \langle a^4 \rangle$. Note that, since $b \notin \langle a \rangle$, we have $bab^{-1}a^3 = e$. Suppose $i, j, k, \ell \in \{\pm 1\}$, with

$$e = \tilde{b}^i \tilde{a}^j \tilde{b}^{-k} \tilde{a}^{3\ell} = \tilde{b}^{i-k} \tilde{a}^{5j+3\ell} \equiv a^{j-\ell} \pmod{\langle \tilde{a}^4 \rangle},$$

so $j = \ell$. Then we must also have $i = k$. We conclude from Lemma B.1 that $\varphi(vs) = \varphi(v)\tilde{s}$, for all $v \in \langle a, b \rangle = G$ and $s \in \{a^{\pm 1}, b^{\pm 1}\}$, so $\varphi \in \text{Aut } G$.

Assume $G = [24, 1]$. Let $a \in S$ with $|a| = 8$, and let $b \in S$, such that $b \notin \langle a \rangle$. Write $\tilde{b} = \tilde{a}^r x$, where $\langle x \rangle = \mathbb{Z}_3$. We may assume $\tilde{b}$ has prime-power order (see Lemma 6.11), and we know that $\tilde{a}^2$ centralizes x , so either r is odd or $r = 0$.

Assume, for the moment, that $r = 0$, which means $\langle \tilde{b} \rangle = \mathbb{Z}_3 = \langle b \rangle$. Then a inverts b , so $aba^{-1}b = e$. Suppose $i, j, k, \ell \in \{\pm 1\}$, with

$$e = \tilde{a}^i \tilde{b}^j \tilde{a}^{-k} \tilde{b}^\ell = \tilde{a}^{i-k} \tilde{b}^{-j+\ell}.$$

Since the exponents of $\tilde{a}$ and $\tilde{b}$ must be 0, we have $i = k$ and $j = \ell$. We conclude from Lemma B.1 that $\varphi(vs) = \varphi(v)\tilde{s}$, for all $v \in \langle a, b \rangle = G$ and $s \in \{a^{\pm 1}, b^{\pm 1}\}$, so $\varphi \in \text{Aut } G$.

We may now assume that r is odd. The proof of Lemma 6.11 shows there is no harm in replacing b with a power that is relatively prime to 8, so we may assume $r = 1$. Since $a^2 \in Z(G)$, we have $a^2ba^{-2}b^{-1} = e$. Suppose $i, j, k, \ell \in \{\pm 1\}$, with

$$e = \tilde{a}^{2i} \tilde{b}^j \tilde{a}^{-2k} \tilde{b}^{-\ell} = \tilde{a}^{2i-2k} \tilde{b}^{j-\ell} \equiv \tilde{a}^{j-\ell} \pmod{\langle \tilde{a}^4, x \rangle}.$$

Then $j = \ell$. Therefore $\tilde{a}^{2i-2k} = e$, so $i = k$. For $v \in G$ with $\varphi(va) = \varphi(v)\tilde{a}$, we conclude from the proof of Lemma B.1 that $\varphi(vba) = \varphi(vb)\tilde{a}$. In addition, interchanging the roles of a and b tells us that if $\varphi(vb) = \varphi(v)\tilde{b}$, then $\varphi(vab) = \varphi(va)\tilde{b}$. We conclude that $\varphi(vs) = \varphi(v)\tilde{s}$, for all $v \in \langle a, b \rangle = G$ and $s \in \{a^{\pm 1}, b^{\pm 1}\}$, so $\varphi \in \text{Aut } G$.

Assume $G = [24, 13]$. We may assume $|s| \in \{2, 3\}$, for all $s \in S$ (see Lemma 6.11). Let $a \in S$ with $|a| = 3$. Choose $b \in S$, such that $b \notin A_4$. Since every element of order 3 is contained in A_4 , we must have $|b| = 2$.

Assume, for the moment, that $\langle a, b \rangle = G$. Note that $(aba^{-1}b)^2 = e$, and, for convenience, let $\tilde{b}_m = \tilde{a}^{-m} \tilde{b} \tilde{a}^m$ for $m \in \mathbb{Z}$. Suppose $i, j, k, \ell \in \{\pm 1\}$, with

$$e = \tilde{a}^i \tilde{b} \tilde{a}^{-j} \tilde{b} \tilde{a}^k \tilde{b} \tilde{a}^{-\ell} \tilde{b} = \tilde{a}^{i-j+k-\ell} \cdot \tilde{b}_{-j+k-\ell} \tilde{b}_{k-\ell} \tilde{b}_{-\ell} \tilde{b}.$$

This implies $k = \ell$, for otherwise 0, $-\ell$, and $k - \ell$ are all distinct modulo 3, so $\tilde{b}_{k-\ell} \tilde{b}_{-\ell} \tilde{b} \equiv \tilde{b}_1 \tilde{b}_{-1} \tilde{b} \equiv e \pmod{\mathbb{Z}_2}$, but $\tilde{b}_{-j+k-\ell}$ is obviously nontrivial $\pmod{\mathbb{Z}_2}$. (Then, since the exponent of $\tilde{a}$ is 0, we must also have $i = j$.) We conclude from Lemma B.1 that $\varphi(vs) = \varphi(v) \tilde{s}$, for all $v \in \langle a, b \rangle = G$ and $s \in \{a^{\pm 1}, b\}$, so $\varphi \in \text{Aut } G$.

We may now assume $\langle a, s \rangle \neq G$, for all $s \in S$. Then, since $b \notin A_4$ (and b is an element of order 2 in S), we see that $b \in Z(G)$. Since $Z(G)$ has only one nontrivial element, this implies that $S = (S \cap A_4) \cup \{b\}$, and that $\tilde{b} = b$ (since only b -edges make 4-cycles with the edges of every other colour). Therefore

$$\text{Cay}(G; S) \cong \text{Cay}(A_4; S \cap A_4) \times \text{Cay}(\mathbb{Z}_2; \{b\}),$$

and $\varphi(b) = b$. Since A_4 is strongly CCA, it is now easy to see that $\varphi \in \text{Aut } G$. $\square$

Example B.3. The following group of order 63 is CCA:

$$G = \mathbb{Z}_7 \rtimes \mathbb{Z}_9 = \langle x, a \mid x^7 = a^9 = e, a^{-1}xa = x^2 \rangle.$$

Proof. Let φ be a colour-preserving automorphism of a connected Cayley graph $\text{Cay}(G; S)$, such that $\varphi(e) = e$. We may assume S is either $\{a^{\pm 1}, x^{\pm 1}\}$ or $\{a^{\pm 1}, (ax)^{\pm 1}\}$, after discarding redundant generators, applying an automorphism of G , and replacing some elements by appropriate powers (cf. the proof of Lemma 6.11).

If $S = \{a^{\pm 1}, x^{\pm 1}\}$, then we may assume $\varphi(x) = x$, by composing with an automorphism of G . Also, since φ is colour-preserving, it must pass to a well-defined automorphism of the cycle $\text{Cay}(G/\langle x \rangle; \{a^{\pm 1}\})$, so there exists $\epsilon \in \{\pm 1\}$, such that $\varphi(ga) = \varphi(g) a^\epsilon$ for all $g \in G$. Then, since $(1, 1)$ is the only pair $(\epsilon, \delta) \in \{\pm 1\}^2$ that satisfies $a^{-\epsilon} x^\delta a^\epsilon = x^2$, we see that $\varphi(x^i a^j) = x^i a^j$ for all i and j , so φ is the identity map, which is certainly affine.

Assume, now, that $S = \{a^{\pm 1}, (ax)^{\pm 1}\}$. Let $a_1 = a$ and $a_2 = xa$. For any $g \in G$ and $\epsilon \in \{\pm 1\}$, if $\varphi(g a_1) = \varphi(g) a_1^\epsilon$, then, since $a_1^3 = a_2^3$ (and φ is colour-preserving), we have $\varphi(g s^m) = \varphi(g) s^{\epsilon m}$, for all m and all $s \in S$. Since S generates G , this implies $\varphi(gs) = \varphi(g) s^\epsilon$ for all g and all $s \in S$. So φ is affine. $\square$